



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/29, A01H 5/00, 5/10	A1	(11) International Publication Number: WO 99/40209 (43) International Publication Date: 12 August 1999 (12.08.99)
(21) International Application Number: PCT/US99/02061 (22) International Filing Date: 27 January 1999 (27.01.99) (30) Priority Data: 09/020,716 9 February 1998 (09.02.98) US (71) Applicant: PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US). (72) Inventors: JUNG, Rudolf; 1549 Germania Drive, Des Moines, IA 50311 (US). BEACH, Larry, R.; 3939 Maquoketa Drive, Des Moines, IA 50311 (US). DRESS, Virginia, M.; 2203 N.W. 90th Street, Clive, IA 50325 (US). RAO, A., Gururaj; 4734 74th Street, Urbandale, IA 50322 (US). RANCH, Jerome, P.; 218 Wilson Road, West Des Moines, IA 50266 (US). ERTL, David, S.; 3325 335th Street, Waukee, IA 50265 (US). HIGGINS, Regina, K.; 1545 44th Street, Des Moines, IA 50311 (US). (74) Agents: MICHEL, Marianne, H. et al.; 7100 N.W. 62nd Avenue, Darwin Building, Johnston, IA 50131-1000 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ALTERATION OF AMINO ACID COMPOSITIONS IN SEEDS (57) Abstract <p>The present invention provides a plant seed the endosperm of which is characterized as having an elevated level of a preselected amino acid. The present invention also provides expression cassettes, vectors, plants, plant cells and a method for enhancing the nutritional value of seeds.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ALTERATION OF AMINO ACID COMPOSITIONS IN SEEDS

BACKGROUND OF THE INVENTION

Feed formulations based on crop plants must typically be supplemented with
5 specific amino acids to provide animals with essential nutrients which are necessary for
their growth. This supplementation is necessary because, in general, crop plants contain
low proportions of several amino acids which are essential for, and cannot be synthesized
by, monogastric animals.

The seeds of crop plants contain different classes of seed proteins. The amino acid
10 composition of these seeds reflects the composition of the prevalent classes of proteins.
Amino acid limitations are usually due to amino acid deficiencies of these prevalent
protein classes.

Among the amino acids necessary for animal nutrition, those that are of limited
availability in crop plants include methionine, lysine, and threonine. Attempts to increase
15 the levels of these amino acids by breeding, mutant selection, and/or changing the
composition of the storage proteins accumulated in the seeds of crop plants, have met with
limited success, or were accompanied by a loss in yield.

For example, although seeds of corn plants containing a mutant transcription factor,
(opaque 2), or a mutant α -zein gene, (floury 2), exhibit elevated levels of total and bound
20 lysine, there is an altered seed endosperm structure which is more susceptible to damage
and pests. Significant yield losses are also typical.

An alternative means to enhance levels of free amino acids in a crop plant is the
modification of amino acid biosynthesis in the plant. The introduction of a feedback-
regulation-insensitive dihydrodipicolinic acid synthase ("DHDPS") gene, which encodes
25 an enzyme that catalyzes the first reaction unique to the lysine biosynthetic pathway, into
plants has resulted in an increase in the levels of free lysine in the leaves and seeds of those
plants. An increase in the levels of free lysine in the embryo results in reduced amount of
oil in the seed. Further free lysine can be lost during the wet milling process reducing the
feed value of the gluten product of the process.

30 The expression of the *lysC* gene, which encodes a mutant bacterial aspartate kinase
that is desensitized to feedback inhibition by lysine and threonine, from a seed-specific
promoter in tobacco plants, has resulted in an increase in methionine and threonine
biosynthesis in the seeds of those plants. See Karchi, *et al.*; The Plant J.; Vol. 3; p. 721;

(1993). However, expression of the *lysC* gene results in only a 6-7% increase in the level of total threonine or methionine in the seed. The expression of the *lysC* gene in seeds has a minimal impact on the nutritional value of those seeds and, thus, supplementation of feed containing *lysC* transgenic seeds with amino acids, such as methionine and threonine, is still required.

There are additional molecular genetic strategies available for enhancing the amino acid quality of plant proteins. Each involves molecular manipulation of plant genes and the generation of transgenic plants.

Protein sequence modification involves the identification of a gene encoding a major protein, preferably a storage protein, as the target for modification to contain more codons of essential amino acids. An important aspect of this approach is to be able to select a region of the protein that can be modified without affecting the overall structure, stability, function, and other cellular and nutritional properties of the protein.

The development of DNA synthesis technology allows the design and synthesis of a gene encoding a new protein with desirable essential amino acid compositions. For example, researchers have synthesized a 292-base pair DNA sequence encoding a polypeptide composed of 80% essential amino acids and used it with the nopaline synthetase (NOS) promoter to construct a chimeric gene. Expression of this gene in the tuber of transgenic potato has resulted in an accumulation of this protein at a level of 0.02% to 0.35% of the total plant protein. This low level accumulation is possibly due to the weak NOS promoter and/or the instability of the new protein.

Tobacco has been used as a test plant to demonstrate the feasibility of this approach by transferring a chimeric gene containing the bean phaseolin promoter and the cDNA of a sulfur-rich protein Brazil Nut Protein ("BNP"), (18 mol% methionine and 8 mol% cysteine) into tobacco. Amino acid analysis indicates that the methionine content in the transgenic seeds is enhanced by 30% over that of the untransformed seeds. This same chimeric gene has also been transferred into a commercial crop, canola, and similar levels of enhancement were achieved.

However, an adverse effect is that lysine content decreases. Additionally, BNP has been identified as a major food allergen. Thus it is neither practical nor desirable to use BNP to enhance the nutritional value of crop plants.

Thus, there is a need to improve the nutritional value of plant seeds. The genetic modification should not be accompanied by detrimental side effects such as allergenicity, anti-nutritional quality or poor yield.

5 **SUMMARY OF THE INVENTION**

It is an object of the present invention to provide a seed, the endosperm of which contains elevated levels of an essential amino acid.

It is a further object of the present invention to provide methods for increasing the nutritional value of feed.

10 It is a further object of the present invention to provide methods for genetically modifying seeds so as to increase amounts of essential amino acids which are present in relatively low amounts in unmodified seeds.

It is a further object of the present invention to provide methods for increasing the nutritional content of seeds without detrimental side effects such as allergenicity or anti-
15 nutritional quality.

It is a further object of the present invention to provide methods for increasing the nutritional content of seeds while maintaining a high yield.

It is a further object of the present invention to provide a method for the expression of a polypeptide in a seed having levels of a preselected amino acid sufficient to reduce or
20 obviate feed supplementation.

According to the present invention a transformed plant seed is provided, the endosperm of which is characterized as having an elevated level of at least one preselected amino acid compared to a seed from a corresponding plant which has not been transformed, wherein the amino acid is lysine, threonine, or tryptophan and optionally a
25 sulfur-containing amino acid.

Also provided is a seed from a plant which has been transformed to express a heterologous protein in the endosperm of the seed, wherein the seed exhibits an elevated level of an essential amino acid.

An expression cassette is also provided comprising a seed endosperm-preferred
30 promoter operably linked to a structural gene encoding a polypeptide having an elevated level of a preselected amino acid. Transformed plants and seeds containing the expression cassette are also provided.

A method for elevating the level of a preselected amino acid in the endosperm of plant seed is also provided. The method comprises the transformation of plant cells by introducing the expression cassette, recovering the transformed cells, regenerating a transformed plant and collecting the seeds therefrom.

5

DETAILED DESCRIPTION OF THE INVENTION

As used herein, a “structural gene ” means an exogenous or recombinant DNA sequence or segment that encodes a polypeptide.

As used herein, “recombinant DNA” is a DNA sequence or segment that has been
10 isolated from a cell, purified, synthesized or amplified.

As used herein, “isolated” means either physically isolated from the cell or synthesized *in vitro* on the basis of the sequence of an isolated DNA segment.

As used herein, the term “increased” or “elevated” levels of the preselected amino acid in a protein means that the protein contains an elevated amount of a preselected amino
15 acid compared to the amount in an average protein.

As used herein, “increased” or “elevated” levels or amounts of preselected amino acids in a transformed plant or seed are levels which are greater than the levels or amounts in the corresponding untransformed plant or seed.

As used herein, “polypeptide” means proteins, protein fragments, modified
20 proteins, amino acid sequences and synthetic amino acid sequences.

As used herein, “transformed plant” means a plant which comprises a structural gene which is introduced into the genome of the plant by transformation.

As used herein, “untransformed plant” refers to a wild type plant, i.e., one where the genome has not been altered by the introduction of the structural gene.

As used herein, “plant” includes but is not limited to plant cells, plant tissue and
25 plant seeds.

As used herein, “seed endosperm-preferred promoter” is a promoter which preferentially promotes expression of the structural gene in the endosperm of the seed.

As used herein with respect to a structural gene encoding a polypeptide, the term
30 “expresses” means that the structural gene is incorporated into the genome of cells, so that the product encoded by the structural gene is produced within the cells.

As used herein, the term "essential amino acid" means an amino acid which is synthesized only by plants or microorganisms or which is not produced by animals in sufficient quantities to support normal growth and development.

As used herein, the term "high lysine content protein" means that the protein has at least about 7 mole % lysine, preferably about 7 mole % to about 50 mole % lysine, more preferably about 7 mole % to about 40 mole % lysine and most preferably about 7 mole % to about 30 mole %.

As used herein, the term "high sulfur content protein" means that the protein contains at least about 6 mole % methionine and/or cysteine, preferably about 6 mole % to about 40 mole %, more preferably about 6 mole % to about 30 mole % and most preferably 6 mole % to 25 mole %.

The present invention provides a transformed plant seed, the endosperm of which is characterized as having an elevated level of a preselected amino acid compared to the seed of a corresponding plant which has not been transformed. It is preferred that the level of preselected amino acid is elevated in the endosperm in preference to other parts of the seed.

The preselected amino acid is an essential amino acid such as lysine, cysteine, methionine, threonine, tryptophan, arginine, valine, leucine, isoleucine, histidine or combinations thereof, preferably, the preselected amino acid is lysine, threonine, cysteine, tryptophan, or combinations thereof and optionally methionine. It is especially preferred that the polypeptide has an increased content of lysine as well as a sulfur containing amino acid, i.e., methionine and/or cysteine.

The polypeptide can be an endogenous or heterologous protein. When an endogenous protein is expressed, the preselected amino acid is lysine, cysteine, threonine, tryptophan, arginine, valine, leucine, isoleucine, histidine or combinations thereof and optionally methionine. When the protein is a heterologous protein, any of the above described preselected amino acids or combinations thereof is present in elevated amounts.

Generally the amount of preselected amino acid in the seed of the present invention is at least about 10 percent by weight greater than in a corresponding untransformed seed, preferably about 10 percent by weight to about 10 times greater, more preferably about 15 percent by weight to about 10 times greater and most preferably about 20 percent to about 10 times greater.

A polypeptide having an elevated amount of the preselected amino acid is expressed in the transformed plant seed endosperm in an amount sufficient to increase the amount of at least one preselected amino acid in the seed of the transformed plant, relative to the amount of the preselected amino acid in the seed of a corresponding untransformed plant.

The choice of the structural gene is based on the desired amino acid composition of the polypeptide encoded by the structural gene, and the ability of the polypeptide to accumulate in seeds. The amino acid composition of the polypeptide can be manipulated by methods, such as site-directed mutagenesis of the structural gene encoding the polypeptide, so as to result in expression of a polypeptide that is increased in the amount of a particular amino acid. For example, site-directed mutagenesis can be used to increase levels of lysine, methionine, cysteine, threonine and/or tryptophan and/or to decrease levels of asparagine and/or glutamine.

The derivatives differ from the wild-type protein by one or more amino acid substitutions, insertions, deletions or the like. Typically, amino acid substitutions are conservative. In the regions of homology to the native sequence, variants preferably have at least 90% amino acid sequence identity, more preferably at least 95% identity.

Typical examples of suitable proteins include barley chymotrypsin inhibitor, barley alpha hordothionin, soybean 2S albumin proteins, rice high methionine protein and sunflower high methionine protein and derivatives of each protein.

Barley alpha hordothionin has been modified to increase the level of particular amino acids. The sequences of genes which express modified alpha hordothionin proteins with enhanced essential amino acids are based on the mRNA sequence of the native *Hordeum vulgare* alpha hordothionin gene (accession number X05901, Ponz *et al.* 1986 Eur. J. Biochem. 156:131-135).

Modified hordothionin proteins are described in U.S. Ser. Nos. 08/838,763 filed April 10, 1997; 08/824,379 filed March 26, 1997; 08/824,382 filed March 26, 1997; and U.S. Pat. No. 5,703,409 issued December 30, 1997 the disclosures of which are incorporated herein in their entirety by reference.

Alpha hordothionin is a 45-amino acid protein which is stabilized by four disulfide bonds resulting from eight cysteine residues. In its native form, the protein is especially rich in arginine and lysine residues, containing 5 residues (10%) of each. However, it is devoid of the essential amino acid methionine.

Alpha hordothionin has been modified to increase the amount of various amino acids such as lysine, threonine or methionine. The protein has been synthesized and the three-dimensional structure determined by computer modeling. The modeling of the protein predicts that the ten charged residues (arginine at positions 5, 10, 17, 19 and 30, and lysine at positions 1, 23, 32, 38 and 45) all occur on the surface of the molecule. The side chains of the polar amino acids (asparagine at position 11, glutamine at position 22 and threonine at position 41) also occur on the surface of the molecule. Furthermore, the hydrophobic amino acids, (such as the side chains of leucine at positions 8, 15, 24 and 33 and valine at position 18) are also solvent- accessible.

The Three-dimensional modeling of the protein indicates that the arginine residue at position 10 is important to retention of the appropriate 3-dimensional structure and possible folding through hydrogen bond interactions with the C-terminal residue of the protein. A lysine, methionine or threonine substitution at that point would disrupt this hydrogen bonding network, leading to a destabilization of the structure. The synthetic peptide having this substitution could not be made to fold correctly, which supported this analysis. Conservation of the arginine residue at position 10 provides a protein which folds correctly.

Alpha hordothionin has been modified to contain 12 lysine residues in the mature hordothionin peptide, referred to as HT12. (Rao *et al.* 1994 Protein Engineering 7(12):1485-1493 and WO 94/16078 published July 21, 1994) The disclosure of each of these is incorporated herein by reference in their entirety.

Further analysis of substitutions which would not alter the 3-dimensional structure of the molecule led to replacement of Asparagine-11, Glutamine-22 and Threonine-41 with lysine residues with virtually no steric hindrance. The resulting compound contains 27% lysine residues.

Other combinations of these substitutions were also made, including changes in amino acid residues at one or more of positions 5, 11, 17, 19, 22, 30 and 41 are lysine, and the remainder of the residues at those positions are the residues at the corresponding positions in the wild type hordothionin.

Since threonine is a polar amino acid, the surface polar amino acid residues, asparagine at position 11 and glutamine at position 22, can be substituted; and the charged amino acids, lysine at positions 1, 23, 32 and 38 and arginine at positions 5, 17, 19, and 30,

can also be substituted with threonine. The molecule can be synthesized by solid phase peptide synthesis.

While the above sequence is illustrative of the present invention, it is not intended to be a limitation. Threonine substitutions can also be performed at positions containing charged amino acids. Only arginine at position 10 and lysine at position 45 are important for maintaining the structure of the protein. One can also substitute at the sites having hydrophobic amino acids. These include positions 8, 15, 18 and 24.

Since methionine is a hydrophobic amino acid, the surface hydrophobic amino acid residues, leucine at positions 8, 15, and 33, and valine at position 18, were substituted with methionine. The surface polar amino acids, asparagine at position 11, glutamine at position 22 and threonine at position 41, are substituted with methionine. The molecule is synthesized by solid phase peptide synthesis and folds into a stable structure. It has seven methionine residues (15.5%) and, including the eight cysteines, the modified protein has a sulfur amino acid content of 33%.

While the above-described proteins are illustrative of suitable polypeptides which can be expressed in the transformed plant, it is not intended to be a limitation. Methionine substitutions can also be performed at positions containing charged amino acids. Only arginine at position 10 is important for maintaining the structure of the protein through a hydrogen-bonding network with serine at position 2 and lysine at position 45. Thus, one can substitute methionine for lysine at positions 1, 23, 32, and/or 38, and for arginine at positions 5, 17, 19 and/or 30.

Many other proteins are also appropriate, for example the protein encoded by the structural gene can be a lysine and/or sulfur rich seed protein, such as the soybean 2S albumin described in U.S. Ser. No. 08/618,911 filed March 20, 1996, and the chymotrypsin inhibitor from barley, Williamson *et al.*, Eur. J Biochem 165: 99-106 (1987), the disclosures of each are incorporated by reference.

Derivatives of these genes can be made by site directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example the gene encoding for the barley high lysine polypeptide (BHL), is derived from barley chymotrypsin inhibitor, U.S. Ser. No. 08/740,682 filed November 1, 1996 and PCT/US97/20441 filed October 31, 1997, the disclosures of each are incorporated herein by reference. The gene encoding for the enhanced soybean albumin gene (ESA), is

derived from soybean 2S albumin described in U.S. Ser. No. 08/618,911, the disclosure of which is incorporated herein in its entirety by reference.

Other examples of sulfur-rich plant proteins within the scope of the invention include plant proteins enriched in cysteine but not methionine, such as the wheat
5 endosperm purothionine (Mak and Jones; Can. J. Biochem.; Vol. 22; p. 83J; (1976); incorporated herein in its entirety by reference), the pea low molecular weight albumins (Higgins, *et al.*; J. Biol. Chem.; Vol. 261; p. 11124; (1986); incorporated herein in its entirety by reference) as well as 2S albumin genes from other organisms. See, for example, Coulter, *et al.*; J. Exp. Bot.; Vol. 41; p. 1541; (1990); incorporated herein in its
10 entirety by reference.

Such proteins also include methionine-rich plant proteins such as from sunflower seed (Lilley, *et al.*; In: Proceedings of the World Congress on Vegetable Protein Utilization in Human Foods and Animal Feedstuffs; Applewhite, H. (ed.); American Oil Chemists Soc.; Champaign, IL; pp. 497-502; (1989); incorporated herein in its entirety by
15 reference), corn (Pedersen, *et al.*; J. Biol. Chem. p. 261; p. 6279; (1986); Kirihaara, *et al.*; Gene, Vol. 71; p. 359; (1988); both incorporated herein in its entirety by reference), and rice (Musumura, *et al.*; Plant Mol. Biol.; Vol. 12; p. 123; (1989); incorporated herein in its entirety by reference).

The present invention also provides a method for genetically modifying plants to
20 increase the level of at least one preselected amino acid in the endosperm of the seed so as to enhance the nutritional value of the seeds.

The method comprises the introduction of an expression cassette into regenerable plant cells to yield transformed plant cells. The expression cassette comprises a seed endosperm-preferred promoter operably linked to a structural gene encoding a polypeptide
25 elevated in content of the preselected amino acid.

A fertile transformed plant is regenerated from the transformed cells, and seeds are isolated from the plant. The structural gene is transmitted through a complete normal sexual cycle of the transformed plant to the next generation.

The polypeptide is synthesized in the endosperm of seed of the plant which has
30 been transformed by insertion of the expression cassette described above. The sequence for the nucleotide molecule, either RNA or DNA, can readily be derived from the amino acid sequence for the selected polypeptide using standard reference texts.

Plants which can be used in the method of the invention include monocotyledonous cereal plants. Preferred plants include maize, wheat, rice, barley, oats, sorghum, millet and rye. The most preferred plant is maize.

Seeds derived from plants regenerated from transformed plant cells, plant parts or plant tissues, or progeny derived from the regenerated transformed plants, may be used directly as feed or food, or further processing may occur.

Transformation

The transformation of plants in accordance with the invention may be carried out in essentially any of the various ways known to those skilled in the art of plant molecular biology. These include, but are not limited to, microprojectile bombardment, microinjection, electroporation of protoplasts or cells comprising partial cell walls, and *Agrobacterium*-mediated DNA transfer.

I. DNA Used for Transformation

DNA useful for introduction into plant cells includes DNA that has been derived or isolated from any source, that may be subsequently characterized as to structure, size and/or function, chemically altered, and later introduced into the plant.

An example of DNA "derived" from a source, would be a DNA sequence or segment that is identified as a useful fragment within a given organism, and which is then synthesized in essentially pure form. An example of such DNA "isolated" from a source would be a useful DNA sequence that is excised or removed from the source by chemical means, e.g., by the use of restriction endonucleases, so that it can be further manipulated, e.g., amplified, for use in the invention, by the methodology of genetic engineering.

Therefore, useful DNA includes completely synthetic DNA, semi-synthetic DNA, DNA isolated from biological sources, and DNA derived from RNA. The DNA isolated from biological sources, or DNA derived from RNA, includes, but is not limited to, DNA or RNA from plant genes, and non-plant genes such as those from bacteria, yeasts, animals or viruses. The DNA or RNA can include modified genes, portions of genes, or chimeric genes, including genes from the same or different genotype.

The term "chimeric gene" or "chimeric DNA" is defined as a gene or DNA sequence or segment comprising at least two DNA sequences or segments from species which do not recombine DNA under natural conditions, or which DNA sequences or

segments are positioned or linked in a manner which does not normally occur in the native genome of untransformed plant.

A structural gene of the invention can be identified by standard methods, e.g., enrichment protocols, or probes, directed to the isolation of particular nucleotide or amino acid sequences. The structural gene can be identified by obtaining and/or screening of a
5 DNA or cDNA library generated from nucleic acid derived from a particular cell type, cell line, primary cells, or tissue.

Screening for DNA fragments that encode all or a portion of the structural gene can be accomplished by screening plaques from a genomic or cDNA library for hybridization
10 to a probe of the structural gene from other organisms or by screening plaques from a cDNA expression library for binding to antibodies that specifically recognize the polypeptide encoded by the structural gene.

DNA fragments that hybridize to a structural gene probe from other organisms and/or plaques carrying DNA fragments that are immunoreactive with antibodies to the polypeptide encoded by the structural gene can be subcloned into a vector and sequenced
15 and/or used as probes to identify other cDNA or genomic sequences encoding all or a portion of the structural gene.

Portions of the genomic copy or copies of the structural gene can be partially sequenced and identified by standard methods including either DNA sequence homology
20 to other homologous genes or by comparison of encoded amino acid sequences to known polypeptide sequences.

Once portions of the structural gene are identified, complete copies of the structural gene can be obtained by standard methods, including cloning or polymerase chain reaction (PCR) synthesis using oligonucleotide primers complementary to the structural gene. The
25 presence of an isolated full-length copy of the structural gene can be verified by comparison of its deduced amino acid sequence with the amino acid sequence of native polypeptide sequences.

As discussed above, the structural gene encoding the polypeptide can be modified to increase the content of particular amino acid residues in that polypeptide by methods
30 well known to the art, including, but not limited to, site-directed mutagenesis. Thus, derivatives of naturally occurring polypeptides can be made by nucleotide substitution of the structural gene so as to result in a polypeptide having a different amino acid at the position in the polypeptide which corresponds to the codon with the nucleotide

substitution. The introduction of multiple amino acid changes in a polypeptide can result in a polypeptide which is significantly enriched in a preselected amino acid.

As noted above, the choice of the polypeptide encoded by the structural gene will be based on the amino acid composition of the polypeptide and its ability to accumulate in seeds. The amino acid can be chosen for its nutritional value to produce a value-added trait to the plant or plant part. Amino acids desirable for value-added traits, as well as a source to limit synthesis of an endogenous protein include, but are not limited to, lysine, threonine, tryptophan, methionine, and cysteine.

10 Expression Cassettes and Expression Vectors

According to the present invention, a structural gene is identified, isolated, and combined with a seed endosperm-preferred promoter to provide a recombinant expression cassette.

The construction of such expression cassettes which can be employed in conjunction with the present invention are well known to those of skill in the art in light of the present disclosure. See, e.g., Sambrook, *et al.*; Molecular Cloning: A Laboratory Manual; Cold Spring Harbor, New York; (1989); Gelvin, *et al.*; Plant Molecular Biology Manual; (1990); Plant Biotechnology: Commercial Prospects and Problems, eds Prakash, *et al.*; Oxford & IBH Publishing Co.; New Delhi, India; (1993); and Heslot, *et al.*; Molecular Biology and Genetic Engineering of Yeasts; CRC Press, Inc., USA; (1992); each incorporated herein in its entirety by reference.

Preferred promoters useful in the practice of the invention are those seed endosperm-preferred promoters that allow expression of the structural gene selectively in seed endosperm to avoid any potential deleterious effects associated with the expression of the structural gene in the embryo.

It has been found that when endosperm-preferred promoters are employed, the total level of the preselected amino acid in the seed is increased compared to a seed produced by employing an embryo-preferred promoter, such as the globulin1 promoter. When the globulin1 promoter is employed, the polypeptide is expressed by the structural gene, but the total amount of the preselected amino acid is not increased.

Examples of suitable promoters include, but are not limited to, 27 kD gamma zein promoter and waxy promoter. See the following sites relating to the 27kD gamma zein promoter: Boronat,A., Martinez,M.C., Reina,M., Puigdomenech,P. and Palau,J.; Isolation

and sequencing of a 28 kD glutelin-2 gene from maize: Common elements in the 5' flanking regions among zein and glutelin genes; Plant Sci. 47, 95-102 (1986) and Reina, M., Ponte, I., Guillen, P., Boronat, A. and Palau, J., Sequence analysis of a genomic clone encoding a Zc2 protein from Zea mays W64 A, Nucleic Acids Res. 18 (21), 6426 (1990). See the following site relating to the waxy promoter: Kloesgen, R.B., Gierl, A., Schwarz-Sommer, ZS. and Saedler, H., Molecular analysis of the waxy locus of Zea mays, Mol. Gen. Genet. 203, 237-244 (1986). The disclosures each of these are incorporated herein by reference in their entirety.

However, other endosperm-preferred promoters can be employed.

II. DELIVERY OF DNA TO CELLS

The expression cassette or vector can be introduced into prokaryotic or eukaryotic cells by currently available methods which are described in the literature. See for example, Weising *et al.*, Ann. Rev. Genet. 2: 421-477 (1988). For example, the expression cassette or vector can be introduced into plant cells by methods including, but not limited to, *Agrobacterium*-mediated transformation, electroporation, PEG poration, microprojectile bombardment, microinjection of plant cell protoplasts or embryogenic callus, silicon fiber delivery, infectious viruses or viroids such as retroviruses, the use of liposomes and the like, all in accordance with well-known procedures.

The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al.*, Embo J. 3: 2717-2722 (1984). Electroporation techniques are described in Fromm *et al.*, Proc. Natl. Acad. Sci. 82: 5324 (1985). Ballistic transformation techniques are described in Klein *et al.*, Nature 327: 70-73 (1987). The disclosure of each of these is incorporated herein in its entirety by reference.

Introduction and expression of foreign genes in plants has been shown to be possible using the T-DNA of the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens*. Using recombinant DNA techniques and bacterial genetics, a wide variety of foreign DNAs can be inserted into T-DNA in *Agrobacterium*. Following infection by the bacterium containing the recombinant Ti plasmid, the foreign DNA is inserted into the host plant chromosomes, thus producing a genetically engineered cell and eventually a genetically engineered plant. A second approach is to introduce root-inducing (Ri) plasmids as the gene vectors.

Agrobacterium tumefaciens-mediated transformation techniques are well described in the literature. See, for example Horsch *et al.*, Science 233: 496-498 (1984), and Fraley *et al.*, Proc. Natl. Acad. Sci. 80: 4803 (1983). *Agrobacterium* transformation of maize is described in U.S. Patent No. 5,550,318. The disclosure of each of these is incorporated
5 herein in its entirety by reference.

Other methods of transfection or transformation include (1) *Agrobacterium rhizogenes*-mediated transformation (see, e.g., Lichtenstein and Fuller In: Genetic Engineering, vol. 6, PWJ Rigby, Ed., London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: DNA Cloning, Vol. II, D. M. Glover, Ed., Oxford, IRI Press, 1985).
10 Application PCT/US87/02512 (WO 88/02405 published Apr. 7, 1988) describes the use of *A. rhizogenes* strain A4 and its Ri plasmid along with *A. tumefaciens* vectors pARC8 or pARC16 (2) liposome-mediated DNA uptake (see, e.g., Freeman *et al.*, Plant Cell Physiol. 25: 1353, 1984), (3) the vortexing method (see, e.g., Kindle, Proc. Natl. Acad. Sci., USA 87: 1228, (1990). The disclosure of each of these is incorporated herein in its entirety by
15 reference.

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou *et al.*, Methods in Enzymology, 101:433 (1983); D. Hess, Intern Rev. Cytol., 107:367 (1987); Luo *et al.*, Plane Mol. Biol. Reporter, 6:165 (1988). The disclosure of each of these is incorporated herein in its entirety by reference.

20 Expression of polypeptide coding genes can be obtained by injection of the DNA into reproductive organs of a plant as described by Pena *et al.*, Nature, 325.:274 (1987). The disclosure of which is incorporated herein in its entirety by reference.

DNA can also be injected directly into the cells of immature embryos and the rehydration of desiccated embryos as described by Neuhaus *et al.*, Theor. Appl. Genet.,
25 75:30 (1987); and Benbrook *et al.*, in Proceedings Bio Expo 1986, Butterworth, Stoneham, Mass., pp. 27-54 (1986). The disclosure of each of these is incorporated herein in its entirety by reference.

Plant cells useful for transformation include cells cultured in suspension cultures, callus, embryos, meristem tissue, pollen, and the like.

30 A variety of plant viruses that can be employed as vectors are known in the art and include cauliflower mosaic virus (CaMV), geminivirus, brome mosaic virus, and tobacco mosaic virus.

Typical vectors useful for expression of genes in higher plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers *et al.*, Meth. In Enzymol., 153:253-277 (1987). These vectors are plant integrating vectors in that on transformation, the vectors integrate a portion of vector DNA into the genome of the host plant. The disclosure of which is incorporated herein in its entirety by reference.

A particularly preferred vector is a plasmid, by which is meant a circular double-stranded DNA molecule which is not a part of the chromosomes of the cell. Exemplary *A. tumefaciens* vectors useful herein are plasmids pKYLX6 and pKYLX7 of Schardl *et al.*, Gene, 61:1-11 (1987) and Berger *et al.*, Proc. Natl. Acad. Sci. U.S.A., 86:8402-8406 (1989). Another useful vector herein is plasmid pBI101.2 that is available from Clontech Laboratories, Inc. (Palo Alto, CA). The disclosure of each of these is incorporated herein in its entirety by reference.

A cell in which the foreign genetic material in a vector is functionally expressed has been "transformed" by the vector and is referred to as a "transformant".

Either genomic DNA or cDNA coding the gene of interest may be used in this invention. The gene of interest may also be constructed partially from a cDNA clone and partially from a genomic clone.

When the gene of interest has been isolated, genetic constructs are made which contain the necessary regulatory sequences to provide for efficient expression of the gene in the host cell.

According to this invention, the genetic construct will contain (a) a genetic sequence coding for the polypeptide of interest and (b) one or more regulatory sequences operably linked on either side of the structural gene of interest. Typically, the regulatory sequences will be a promoter or a terminator. The regulatory sequences may be from autologous or heterologous sources.

The cloning vector will typically carry a replication origin, as well as specific genes that are capable of providing phenotypic selection markers in transformed host cells. Typically, genes conferring resistance to antibiotics or selected herbicides are used. After the genetic material is introduced into the target cells, successfully transformed cells and/or colonies of cells can be isolated by selection on the basis of these markers.

Typical selectable markers include genes coding for resistance to the antibiotic spectinomycin (e.g., the *aada* gene), the streptomycin phosphotransferase (SPT) gene

coding for streptomycin resistance, the neomycin phosphotransferase (NPTII) gene encoding kanamycin or geneticin resistance, the hygromycin phosphotransferase (HPT) gene coding for hygromycin resistance.

Genes coding for resistance to herbicides include genes which act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonylurea-type herbicides (e.g., the acetolactate synthase (ALS) genes containing mutations leading to such resistance in particular the S4 and/or Hra mutations), genes coding for resistance to herbicides which act to inhibit action of glutamine synthase, such as phosphinothricin or basta (e.g., the *pat* or *bar* gene), or other such genes known in the art. The *bar* gene encodes resistance to the herbicide basta, and the ALS gene encodes resistance to the herbicide chlorsulfuron.

Typically, an intermediate host cell will be used in the practice of this invention to increase the copy number of the cloning vector. With an increased copy number, the vector containing the gene of interest can be isolated in significant quantities for introduction into the desired plant cells.

Host cells that can be used in the practice of this invention include prokaryotes, including bacterial hosts such as *E. coli*, *S. typhimurium*, and *Serratia marcescens*. Eukaryotic hosts such as yeast or filamentous fungi may also be used in this invention. Since these hosts are also microorganisms, it will be essential to ensure that plant promoters which do not cause expression of the polypeptide in bacteria are used in the vector.

The isolated cloning vector will then be introduced into the plant cell using any convenient transformation technique as described above.

III. Regeneration and Analysis of Transformants

Following transformation, regeneration is involved to obtain a whole plant from transformed cells and the presence of structural gene (s) or "transgene(s)" in the regenerated plant is detected by assays. The seed derived from the plant is then tested for levels of preselected amino acids. Depending on the type of plant and the level of gene expression, introduction of the structural gene into the plant seed endosperm can enhance the level of preselected amino acids in an amount useful to supplement the nutritional quality of those seeds.

Using known techniques, protoplasts and cell or tissue culture can be regenerated to form whole fertile plants which carry and express the gene for a polypeptide according to this invention.

Accordingly, a highly preferred embodiment of the present invention is a transformed maize plant, the cells of which contain at least one copy of the DNA sequence of an expression cassette containing a gene encoding a polypeptide containing elevated amounts of an essential amino acid, such as HT12, BHL or ESA protein.

Techniques for regenerating plants from tissue culture, such as transformed protoplasts or callus cell lines, are known in the art. For example, see Phillips, *et al.*; Plant Cell Tissue Organ Culture; Vol. 1; p. 123; (1981); Patterson, *et al.*; Plant Sci.; Vol. 42; p. 125; (1985); Wright, *et al.*; Plant Cell Reports; Vol. 6; p. 83; (1987); and Barwale, *et al.*; Planta; Vol. 167; p. 473; (1986); each incorporated herein in its entirety by reference. The selection of an appropriate method is within the skill of the art.

Examples of the practice of the present invention detailed herein relate specifically to maize plants. However, the present invention is also applicable to other cereal plants. The expression vectors utilized herein are demonstrably capable of operation in cells of cereal plants both in tissue culture and in whole plants. The invention disclosed herein is thus operable in monocotyledonous species to transform individual plant cells and to achieve full, intact plants which can be regenerated from transformed plant cells and which express preselected polypeptides.

The introduced structural genes are expressed in the transformed plant cells and stably transmitted (somatically and sexually) to the next generation of cells produced. The vector should be capable of introducing, maintaining, and expressing a structural gene in plant cells. The structural gene is passed on to progeny by normal sexual transmission.

To confirm the presence of the structural gene (s) or "transgene(s)" in the regenerating plants, or seeds or progeny derived from the regenerated plant, a variety of assays can be performed. Such assays include Southern and Northern blotting; PCR; assays that detect the presence of a polypeptide product, e.g., by immunological means (ELISAs and Western blots) or by enzymatic function; plant part assays, such as leaf, seed or root assays; and also, by analyzing the phenotype of the whole regenerated plant.

Whereas DNA analysis techniques can be conducted using DNA isolated from any part of a plant, RNA will be expressed in the seed endosperm and hence it will be necessary to prepare RNA for analysis from these tissues.

PCR techniques can be used for detection and quantitation of RNA produced from introduced structural genes. In this application of PCR it is first necessary to reverse transcribe RNA into DNA, using enzymes such as reverse transcriptase, and then through the use of conventional PCR techniques amplify the DNA. In most instances PCR techniques, while useful, will not demonstrate integrity of the RNA product.

Further information about the nature of the RNA product may be obtained by Northern blotting. This technique will demonstrate the presence of an RNA species and give information about the integrity of that RNA. The presence or absence of an RNA species can also be determined using dot or slot blot Northern hybridizations. These techniques are modifications of Northern blotting and will only demonstrate the presence or absence of an RNA species.

While Southern blotting and PCR may be used to detect the structural gene in question, they do not provide information as to whether the structural gene is being expressed. Expression may be evaluated by specifically identifying the polypeptide products of the introduced structural genes or evaluating the phenotypic changes brought about by their expression.

Assays for the production and identification of specific polypeptides may make use of physical-chemical, structural, functional, or other properties of the polypeptides. Unique physical-chemical or structural properties allow the polypeptides to be separated and identified by electrophoretic procedures, such as native or denaturing gel electrophoresis or isoelectric focusing, or by chromatographic techniques such as ion exchange or gel exclusion chromatography.

The unique structures of individual polypeptides offer opportunities for use of specific antibodies to detect their presence in formats such as an ELISA assay.

Combinations of approaches may be employed with even greater specificity such as Western blotting in which antibodies are used to locate individual gene products that have been separated by electrophoretic techniques.

Additional techniques may be employed to absolutely confirm the identity of the product of interest such as evaluation by amino acid sequencing following purification.

Although these are among the most commonly employed, other procedures may be additionally used.

Very frequently, the expression of a gene product is determined by evaluating the phenotypic results of its expression. These assays also may take many forms, including

but not limited to, analyzing changes in the chemical composition, morphology, or physiological properties of the plant. In particular, the elevated preselected amino acid content due to the expression of structural genes encoding polypeptides can be detected by amino acid analysis.

5 Breeding techniques useful in the present invention are well known in the art.

The present invention will be further described by reference to the following detailed examples. It is understood, however, that there are many extensions, variations, and modifications on the basic theme of the present invention beyond that shown in the examples and description, which are within the spirit and scope of the present invention.

10

Examples

EXAMPLE 1

Construction of the HT12 gene and of other genes encoding polypeptides having an elevated level of a preselected amino acid.

15

As noted above, the sequence of the HT12 gene is based on the mRNA sequence of the native *Hordeum vulgare* alpha hordothionin gene (accession number X05901, Ponz *et al.* 1986 Eur. J. Biochem. 156:131-135) modified to introduce 12 lysine residues into the mature hordothionin peptide (See Rao *et al.* 1994 Protein Engineering 7(12):1485-1493, and WO 94/16078 published July 21, 1994).

20

The alpha hordothionin cDNA comprising the entire alpha hordothionin coding sequence is isolated by rt-PCR of mRNA from developing barley seed. Primers are designed based upon the published alpha hordothionin sequence to amplify the gene and to introduce a NcoI site at the start (ATG) codon and a BamHI site after the stop codon of the thionin coding sequence to facilitate cloning.

25

Primers are designated as HTPCR1 (5'-AGTATAAGTAAACACACCATCACACCCTTGAGGCCCTTGCTGGTGGCCATGGT G-3') and HTPCR2 (5'-CCTCACATCCCTTAGTGCCTAAGTTCGACGTCGGGCCCTCTAGTCGACGGATCC A-3'). These primers are used in a PCR reaction to amplify alpha hordothionin by conventional methods. The resulting PCR product is purified and subcloned into the BamHI/NcoI digested pBSKP vector (Stratagene, LaJolla, CA) and sequenced on both strands to confirm its identity. The clone is designated pBSKP-HT (seq. ID 1). Primers are designed for single stranded DNA site-directed mutagenesis to introduce 12 codons for

30

lysine, based on the peptide structure of hordothionin 12 (Ref: Rao *et al.* 1994 Protein Engineering 7(12):1485-1493) and are designated HT12mut1 (5'-AGCGGAAAATGCCCCGAAAGGCTTCCCCAAATTGGC-3'), HT12mut2 (5'-TGCGCAGGCGTCTGCAAGTGTAAGCTGACTAGTAGCGGAAAATGC-3'),
5 HT12mut3 (5'-TACAACCTTTGCAAAGTCAAAGGCGCCAAGAAGCTTTGCGCAGGCGTCTG-3'),
HT12mut4 (5'-GCAAGAGTTGCTGCAAGAGTACCCTGGGAAGGAAGTGCTACAACCTTTGC-3').

Sequence analysis is used to verify the desired sequence of the resulting plasmid,
10 designated pBSKP-HT12 (seq. ID 2).

Similarly, genes encoding other derivatives of hordothionine, as described above, (See U.S. Ser. Nos. 08/838,763 filed April 10, 1997; 08/824,379 filed March 26, 1997; 08/824,382 filed March 26, 1997; and U.S. Pat. No. 5,703,409 issued December 30, 1997), the gene encoding enhanced soybean albumin (ESA) (See U.S. Ser. No. 08/618,911), and
15 genes encoding BHL and other derivatives of the barley chymotrypsin inhibitor (See U.S. Ser. No. 08/740,682 filed November 1, 1996 and PCT/US97/20441 filed October 31, 1997) are constructed by site directed mutagenesis from pBSKP-HT, a subclone of the soybean 2S albumin 3 gene in the pBSKP vector (Stratagene, LaJolla, CA), and a subclone of the barley chymotrypsin inhibitor in the pBSKP vector, respectively.

20

EXAMPLE 2

Construction of vectors for seed preferred expression of polypeptides having an elevated level of a preselected amino acid.

25 A 442bp DNA fragment containing the modified hordothionin gene encoding HT12 is isolated from plasmid pBSKP-HT12 by NcoI/BamHI restriction digestion, gel purification and is ligated between the 27 kD gamma zein promoter and 27kD gamma zein terminator of the NcoI/BamHI digested vector PHP3630. PHP 3630 is a subclone of the endosperm-preferred 27kD gamma zein gene (Genbank accession number X58197) in the
30 pBSKP vector (Stratagene), which is modified by site directed mutagenesis by insertion of a NcoI site at the start codon (ATG) of the 27kD gamma zein coding sequence. The 27kD gamma zein coding sequence is replaced with the HT12 coding sequence. The resulting expression vector containing the chimeric gene construct gz::HT12::gz, designated as

PHP8001 (Seq. ID 3), is verified by extensive restriction digest analysis and DNA sequencing.

Similarly, the 442bp DNA fragment containing the HT12 coding sequence is inserted between the globulin1 promoter and the globulin1 terminator of the embryo preferred corn globulin1 gene (Genbank accession number X59083), and between the waxy promoter and the waxy terminator of the endosperm-preferred waxy gene (Genbank accession number M24258). The globulin1 and waxy coding sequences, respectively, are replaced with the HT12 coding sequence. The resulting chimeric genes *glb1::HT12::glb1*, and *wx::HT12::wx* are designated as PHP 7999 (Seq. ID 4), and PHP 5025 (Seq. ID 5).

In a like manner, expression vectors containing genes encoding other derivatives of hordothionine (See Rao *et al.* 1994 Protein Engineering 7(12):1485-1493, and WO 94/16078 published July 21, 1994), the gene encoding enhanced soybean albumin (ESA) (See U.S. Ser. No. 08/618,911), and genes encoding BHL and other derivatives of the barley chymotrypsin inhibitor (See U.S. Ser. No. 08/740,682 filed November 1, 1996 and PCT/US97/20441 filed October 31, 1997) are constructed by insertion of the corresponding coding sequences between the promoter and terminator of the 27kD gamma zein gene, the globulin1 gene and the waxy gene, respectively. Resulting chimeric genes are for example *gz::ESA::gz* and *gz::BHL::gz*, designated as PHP11260 (Seq. ID 6) and as PHP11427 (Seq. ID 7), respectively.

The resulting expression vectors are used in conjunction with the selectable marker expression cassettes PHP3528 (enhanced CAMV::Bar::PinII) for particle bombardment transformation of maize immature embryos.

EXAMPLE 3

Preparation of Transgenic Plants

The general method of genetic transformation used to produce transgenic maize plants is mediated by bombardment of embryogenically responsive immature embryos with tungsten particles associated with DNA plasmids, said plasmids consisting of a selectable and an unselectable marker gene.

Preparation of Tissue

Immature embryos of "High Type II" are the target for particle bombardment-mediated transformation. This genotype is the F₁ of two purebred genetic lines, parent A

and parent B, derived from A188 X B73. Both parents are selected for high competence of somatic embryogenesis. See Armstrong, *et al.*, "Development and Availability of Germplasm with High Type II Culture Formation Response," Maize Genetics Cooperation Newsletter, Vol. 65, pp. 92 (1991); incorporated herein in its entirety by reference.

5 Ears from F₁ plants are selfed or sibbed, and embryos are aseptically dissected from developing caryopses when the scutellum first becomes opaque. The proper stage occurs about 9-13 days post-pollination, and most generally about 10 days post-pollination, and depends on growth conditions. The embryos are about 0.75 to 1.5 mm long. Ears are surface sterilized with 20-50% Clorox for 30 min, followed by 3 rinses with sterile
10 distilled water.

Immature embryos are cultured, scutellum oriented upward, on embryogenic induction medium comprised of N6 basal salts (Chu, *et al.*, "Establishment of an Efficient Medium for Anther Culture of Rice Through Comparative Experiments on the Nitrogen Sources," Scientia Sinica, (Peking), Vol. 18, pp. 659-668 (1975); incorporated herein in its
15 entirety by reference; Eriksson vitamins (See Eriksson, T., "Studies on the Growth Requirements and Growth Measurements of Haplopappus gracilis," Physiol. Plant, Vol. 18, pp. 976-993 (1965); incorporated herein in its entirety by reference), 0.5 mg/l thiamine HCl, 30 gm/l sucrose, 2.88 gm/l L-proline, 1 mg/l 2,4-dichlorophenoxyacetic acid, 2 gm/l Gelrite, and 8.5 mg/l AgNO₃.

20 The medium is sterilized by autoclaving at 121°C for 15 min and dispensed into 100 X 25 mm petri dishes. AgNO₃ is filter-sterilized and added to the medium after autoclaving. The tissues are cultured in complete darkness at 28°C. After about 3 to 7 days, generally about 4 days, the scutellum of the embryo has swelled to about double its original size and the protuberances at the coleorhizal surface of the scutellum indicate the
25 inception of embryogenic tissue. Up to 100% of the embryos display this response, but most commonly, the embryogenic response frequency is about 80%.

When the embryogenic response is observed, the embryos are transferred to a medium comprised of induction medium modified to contain 120 gm/l sucrose. The embryos are oriented with the coleorhizal pole, the embryogenically responsive tissue,
30 upwards from the culture medium. Ten embryos per petri dish are located in the center of a petri dish in an area about 2 cm in diameter. The embryos are maintained on this medium for 3-16 hr, preferably 4 hours, in complete darkness at 28°C just prior to

bombardment with particles associated with plasmid DNAs containing the selectable and unselectable marker genes.

To effect particle bombardment of embryos, the particle-DNA agglomerates are accelerated using a DuPont PDS-1000 particle acceleration device. The particle-DNA agglomeration is briefly sonicated and 10 μ l are deposited on macrocarriers and the ethanol allowed to evaporate. The macrocarrier is accelerated onto a stainless-steel stopping screen by the rupture of a polymer diaphragm (rupture disk). Rupture is effected by pressurized helium. Depending on the rupture disk breaking pressure, the velocity of particle-DNA acceleration may be varied. Rupture disk pressures of 200 to 1800 psi are commonly used, with those of 650 to 1100 psi being more preferred, and about 900 psi being most highly preferred. Rupture disk breaking pressures are additive so multiple disks may be used to effect a range of rupture pressures.

Preferably, the shelf containing the plate with embryos is 5.1 cm below the bottom of the macrocarrier platform (shelf #3), but may be located at other distances. To effect particle bombardment of cultured immature embryos, a rupture disk and a macrocarrier with dried particle-DNA agglomerates are installed in the device. The He pressure delivered to the device is adjusted to 200 psi above the rupture disk breaking pressure. A petri dish with the target embryos is placed into the vacuum chamber and located in the projected path of accelerated particles. A vacuum is created in the chamber, preferably about 28 inches Hg. After operation of the device, the vacuum is released and the petri dish is removed.

Bombarded embryos remain on the osmotically adjusted medium during bombardment, and preferably for two days subsequently, although the embryos may remain on this medium for 1 to 4 days. The embryos are transferred to selection medium comprised of N6 basal salts, Eriksson vitamins, 0.5 mg/l thiamine HCl, 30 gm/l sucrose, 1 mg/l 2,4-dichlorophenoxyacetic acid, 2 gm/l Gelrite, 0.85 mg/l AgNO₃ and 3 mg/l bialaphos. Bialaphos is added filter-sterilized. The embryos are subcultured to fresh selection medium at 10 to 14 day intervals. After about 7 weeks, embryogenic tissue, putatively transgenic for both selectable and unselected marker genes, is seen to proliferate from about 7% of the bombarded embryos. Putative transgenic tissue is rescued, and that tissue derived from individual embryos is considered to be an event and is propagated independently on selection medium. Two cycles of clonal propagation is achieved by visual selection for the smallest contiguous fragments of organized embryogenic tissue.

For regeneration of transgenic plants, embryogenic tissue is subcultured to medium comprised of MS salts and vitamins (Murashige, T. and F. Skoog, "A revised medium for rapid growth and bio assays with tobacco tissue cultures"; Physiologia Plantarum; Vol. 15; pp. 473-497; 1962; incorporated herein in its entirety by reference), 100 mg/l myo-inositol, 60 gm/l sucrose, 3 gm/l Gelrite, 0.5 mg/l zeatin, 1 mg/l indole-3-acetic acid, 26.4 ng/l cis-trans-abscissic acid, and 3 mg/l bialaphos in 100 X 25 mm petri dishes and incubated in darkness at 28°C until the development of well-formed, matured somatic embryos can be visualized. This requires about 14 days.

Well-formed somatic embryos are opaque and cream-colored, and are comprised of an identifiable scutellum and coleoptile. The embryos are individually subcultured to germination medium comprised of MS salts and vitamins, 100 mg/l myo-inositol, 40 gm/l sucrose and 1.5 gm/l Gelrite in 100 X 25 mm petri dishes and incubated under a 16 hr light: 8 hr dark photoperiod and 40 $\mu\text{Einsteinsm}^{-2}\text{sec}^{-1}$ from cool-white fluorescent tubes. After about 7 days, the somatic embryos have germinated and produced a well-defined shoot and root. The individual plants are subcultured to germination medium in 125 x 25 mm glass tubes to allow further plant development. The plants are maintained under a 16 hr light: 8 hr dark photoperiod and 40 $\mu\text{Einsteinsm}^{-2}\text{sec}^{-1}$ from cool-white fluorescent tubes.

After about 7 days, the plants are well-established and are transplanted to horticultural soil, hardened off, and potted into commercial greenhouse soil mixture and grown to sexual maturity in a greenhouse. An elite inbred line is used as a male to pollinate regenerated transgenic plants.

Preparation of Particles

Fifteen mg of tungsten particles (General Electric) , 0.5 to 1.8 μm , preferably 1 to 1.8 μm , and most preferably 1 μm , are added to 2 ml of concentrated nitric acid. This suspension is sonicated at 0°C for 20 min (Branson Sonifier Model 450, 40% output, constant duty cycle). Tungsten particles are pelleted by centrifugation at 10,000 rpm (Biofuge) for 1 min and the supernatant is removed. Two ml of sterile distilled water is added to the pellet and sonicate briefly to resuspend the particles. The suspension is pelleted, 1 ml of absolute ethanol is added to the pellet and sonicated briefly to resuspend the particles. Rinse, pellet, and resuspend the particles a further 2 times with sterile

distilled water, and finally resuspend the particles in 2 ml of sterile distilled water. The particles are subdivided into 250 µl aliquots and stored frozen.

Preparation of particle-plasmid DNA association

- 5 The stock of tungsten particles is sonicated briefly in a water bath sonicator (Branson Sonifier Model 450, 20% output, constant duty cycle) and 50 µl is transferred to a microfuge tube. Plasmid DNA is added to the particles for a final DNA amount of 0.1 to 10 µg in 10 µl total volume, and briefly sonicated. Preferably 1 µg total DNA is used. Specifically, 5 µl of PHP8001 (gz::HT12::gz) and 5µl of PHP3528 (enhanced
- 10 CAMV::Bar::PinII), at 0.1 µg/µl in TE buffer, are added to the particle suspension. Fifty µl of sterile aqueous 2.5 M CaCl₂ are added, and the mixture is briefly sonicated and vortexed. Twenty µl of sterile aqueous 0.1M spermidine are added and the mixture is briefly sonicated and vortexed. The mixture is incubated at room temperature for 20 min with intermittent brief sonication. The particle suspension is centrifuged, and the
- 15 supernatant is removed. Two hundred fifty µl of absolute ethanol is added to the pellet and briefly sonicated. The suspension is pelleted, the supernatant is removed, and 60 µl of absolute ethanol is added. The suspension is sonicated briefly before loading the particle-DNA agglomeration onto macrocarriers.

20 EXAMPLE 4

Analysis of seed from transgenic plants for recombinant polypeptides having an elevated level of a preselected amino acid.

Preparation of meals from corn seed

- 25 Pooled or individual dry seed harvested from transformed plants from the greenhouse or the field are prepared in one of the following ways:
- A. Seed is imbibed in sterile water overnight (16-20 hr) at 4°C. The imbibed seed is dissected into embryo, endosperm and pericarp. The embryos and endosperm are separately frozen in liquid N₂, the pericarps are discarded. Frozen tissue is ground
- 30 with a liquid N₂ chilled ceramic mortar and pestle to a fine meal. The meals are dried under vacuum and stored at -20°C or -80°C.
- B. Dry whole seed is ground to a fine meal with a ball mill (Klecko), or by hand with a ceramic mortar and pestle. For analysis of endosperm only, the embryos are

removed with a drill and discarded. The remaining endosperm with pericarp is ground with a ball mill or a mortar and pestle.

ELISA analysis

5 Rabbit polyclonal anti HT12 antisera are produced with synthetic HT12 (See Rao *et al. supra*) at Bethyl laboratories. An HT12 ELISA assay is developed and performed by the Analytical Biochemistry department of Pioneer Hi-Bred International, Inc., essentially as described by Harlow and Lane, Antibodies, A Laboratory Manual, Cold Springs Harbor Publication, New York (1988). Quantitative ELISA assays are first performed on pooled
10 meals to identify positive events. Positive events are further analyzed by quantitative ELISA on individual kernels to determine the relative level of HT12 expression and transgene segregation ratio. Among 97 events tested, 59 show HT12 expression levels >1000 ppm. The highest events have HT12 expression levels at 2-5% of the total seed protein. Typical results for HT12 levels for whole kernels of wild type corn, for one event (TC2031) of corn
15 transformed with the *gz::HT12::gz* chimeric gene, expressing HT12 in the endosperm, for one event (TC320) of corn transformed with the *wx::HT12::wx* chimeric gene, expressing HT12 in the endosperm, and for one event (TC2027) of corn transformed with the *glb1::HT12::glb1* chimeric gene, expressing HT12 in the embryo, are in Table 1.

Similarly, antisera are produced, ELISA assays are developed and assays of seed from
20 transformed plants are performed for other derivatives of hordothionine (See Rao *et al. supra* 1994 Protein Engineering 7(12):1485-1493, and WO 94/16078 published July 21, 1994), for the enhanced soybean albumin (ESA) (See U.S. Ser. No. 08/618,911) and for BHL and other derivatives of the barley chymotrypsin inhibitor (See U.S. Ser. No. 08/740,682 filed November 1, 1996 and PCT/US97/20441 filed October 31, 1997), respectively.

25

Polyacrylamide gel and immuno blot analysis

SDS extracts of meals, molecular weight markers, and a synthetic HT12 positive control (see Rao *et al. supra*) are separated on 16.5% or 8-22% polyacrylamide gradient Tris-Tricine gels (Schagger, H. and Von Jagow, G. 1987 Anal. Biochem., 166:368). For immuno
30 blot analysis, gels are transferred to PVDF membranes in 100 mM CAPS, pH 11; 10% methanol using a semidry blotter (Hoefer, San Francisco, CA). After transfer the membrane is blocked in BLOTTO (4% dry milk in Tris-buffered saline, pH 7.5) (Johnson, D. A. ,

Gausch, J. W., Sportsman, J. R., and Elder, J. H. 1984, Gene Anal. Techn., 1:3). The blots are incubated with rabbit anti-HT12 (same as used for ELISA) diluted 1:2000 to 1:7500 in BLOTTO 2 hr at room temperature (22°C) or overnight at 4°C. Blots are washed 4-5X with BLOTTO, then incubated 1-2 hr with horseradish peroxidase-goat anti-rabbit IgG (Promega, Madison, WI) diluted 1:7500 to 1:15000 in BLOTTO. After secondary antibody, the blots are washed 3X with BLOTTO followed by 2 washes with Tris-buffered saline, pH 7.5. Blots are briefly incubated with enhanced chemiluminescence (ECL, Amersham, Arlington Heights, IL) substrate, and wrapped in plastic wrap. Reactive bands are visualized after exposure to x-ray film (Kodak Biomax MR) after short exposure times ranging from 5-120 sec.

HT12 transgenic seed shows a distinctive band not seen in wild type seed at the correct molecular weight and position as judged by the HT12 positive control standard and molecular weight markers. These results indicate that the expressed HT12 prepropeptide is being correctly processed like native HT in barley. Novel polypeptide bands co-migrating with the HT12 positive control are also observed in Coomassie stained polyacrylamide gels loaded with 10mg total extracted protein indicating substantial expression and accumulation of HT12 protein in the seed.

Similarly, other derivatives of hordothionin, soybean albumin, the enhanced soybean albumin (ESA), BHL and other derivatives of the barley chymotrypsin inhibitor are detected by polyacrylamide gel and immuno blot analysis.

Amino acid composition analysis

Meals from seed, endosperm or embryo that express a recombinant polypeptide having an elevated level of a preselected amino acid are sent to the University of Iowa Protein Structure Facility for amino acid composition analysis using standard protocols for digestion and analysis.

Typical results for the amino acid composition of whole kernels of wild type corn, for one event (TC2031) of corn transformed with the gz::HT12::gz chimeric gene, expressing HT12 in the endosperm, for one event (TC320) of corn transformed with the wx::HT12::wx chimeric gene, expressing HT12 in the endosperm, and for one event (TC2027) of corn transformed with the glb1::HT12::glb1 chimeric gene, expressing HT12 in the embryo, are in Table 1.

Table 1: HT12 ELISA analysis and amino acid composition of meal from whole kernels from wild type corn and from transformed corn expressing recombinant HT12.—

transgene	none	wx::HT12::wx	gz::HT12::gz	glb1::HT12::glb1
event	wild-type	TC320	TC2031	TC2027
ELISA				
HT 12	protein ppm 0.00	protein ppm 6200	protein ppm 8000	protein ppm 22600
AA				
	Meal % n=3	Meal % n=2	Meal % n=3	Meal % n=4
Lys	0.29	0.38	0.39	0.24
Arg	0.52	0.58	0.56	0.45
Cys	0.12	0.19	0.17	0.22

- 5 The results in Table 1 demonstrate corn expressing recombinant HT12 in the endosperm shows a significant increase of the preselected amino acid lysine.

Table 2: SEQUENCE INFORMATION

SEQUENCE ID	PROMOTER	GENE
Seq. 1: pBSKP-HT	None	3361-2947
Seq. 2: pBSKP-HT12	None	3361-2947
Seq. 3: PHP8001gz::HT12::gz expression vector	676-2198	2199-2612
Seq. 4: PHP7999 glb1::HT12::glb1 expression vector	3271-1834	1834-1420
Seq. 5: PHP5025 wx::HT::wx expression vector	43-1342	1343-1757
Seq. 6: PHP 11260 gz::ESA::gz expression vector	676-2198	2199-2675
Seq. 7: PHP11427 gz::BHL::gz	676-2198	2199-2450

10

The invention is not limited to the exact details shown and described, for it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention defined by the claims.

WHAT IS CLAIMED IS:

1. A transformed cereal plant seed, the endosperm of which is characterized as having an elevated level of at least one preselected amino acid compared to a seed from a corresponding plant which has not been transformed, wherein the amino
5 acid is lysine, cysteine, threonine, tryptophan, arginine, valine, leucine, isoleucine, histidine or combinations thereof and optionally methionine.
2. The seed according to claim 1 wherein the preselected amino acid is lysine, threonine or tryptophan and optionally a sulfur-containing amino acid.
3. The seed according to Claim 2 wherein the preselected amino acid is lysine.
- 10 4. The seed according to Claim 3 wherein the preselected amino acid is lysine and a sulfur-containing amino acid.
5. The seed according to Claim 1 wherein the plant is selected from the group consisting of maize, wheat, rice, barley, oats, sorghum, millet and rye.
6. The seed according to Claim 5 which is a maize seed.
- 15 7. The seed according to Claim 1 wherein the plant expresses a transgenic protein having an elevated level of the preselected amino acid.
8. The seed according to Claim 7 wherein the protein is barley chymotrypsin inhibitor, barley alpha hordothionin, soybean 2S albumin protein, rice high methionine protein, sunflower high methionine protein or derivatives of each
20 protein.
9. The seed according to Claim 1 wherein the amount of preselected amino acid in the seed is increased at least about 10 percent by weight compared to a corresponding seed which has not been transformed.
10. The seed according to Claim 9 wherein the amount of the preselected amino acid in
25 the seed is about 10 percent by weight to about 10 times greater compared to a corresponding seed which has not been transformed.
11. The seed according to Claim 10 wherein the amount of the preselected amino acid in the seed is about 15 percent by weight to about 10 times greater compared to a corresponding seed which has not been transformed.
- 30 12. The seed according to Claim 11 wherein the amount of the preselected amino acid in the seed is about 20 percent by weight to about 10 times greater compared to a corresponding seed which has not been transformed.

13. An expression cassette comprising a seed endosperm-preferred promoter operably linked to a structural gene encoding a polypeptide elevated in content of a preselected amino acid.
14. The cassette according to Claim 13 wherein the promoter is a gamma zein promoter or a waxy promoter.
15. A vector comprising the expression cassette of Claim 13.
16. A plant cell transformed with the vector of Claim 15.
17. A transformed plant comprising the vector of Claim 15.
18. A seed product obtainable from the transformed seed of Claim 1.
19. A seed from a cereal plant which has been transformed to express a heterologous protein in the endosperm of the seed, wherein the seed exhibits an elevated level of an essential amino acid compared to a plant which has not been transformed.
20. A method for increasing the nutritional value of a cereal plant seed comprising: transforming a host plant cell with a vector comprising an expression cassette comprising a seed endosperm-preferred promoter operably linked to a structural gene encoding a polypeptide elevated in content of a preselected amino acid; recovering the transformed cells; regenerating a transformed plant; and recovering the seeds therefrom.
21. A seed produced by the method of claim 20.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- 5 (i) APPLICANT: Jung, Rudolf
Beach, Larry R.
Dress, Virginia M.
Rao, A. Gururaj
10 Ranch, Jerome P.
Ertl, David S.
Higgins, Regina K.
- (ii) TITLE OF THE INVENTION: Alteration of Amino Acid Compositions
in Seeds
- 15 (iii) NUMBER OF SEQUENCES: 13
- (iv) CORRESPONDENCE ADDRESS:
20 (A) ADDRESSEE: Pioneer Hi-Bred International, Inc.
(B) STREET: 7100 NW 62nd Avenue, P.O. Box 1000
(C) CITY: Johnston
(D) STATE: IA
(E) COUNTRY: USA
(F) ZIP: 50131
- 25 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
30 (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
35 (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
40
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Michel, Marianne H
45 (B) REGISTRATION NUMBER: 35,286
(C) REFERENCE/DOCKET NUMBER: 0815
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 515-334-4467
50 (B) TELEFAX: 515-334-6883
(C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
- 55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 3363 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
60 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
65 TCGACCTCGA GGGGGGGCCC GGTACCCAGC TTTTGTCC TTTAGTGAGG GTTAATTGCG 60

CGCTTGGCGT AATCATGGTC ATAGCTGTTT CCTGTGTGAA ATTGTTATCC GCTCACAATT 120
CCACACAACA TACGAGCCGG AAGCATAAAG TGTAAGCCT GGGGTGCCTA ATGAGTGAGC 180
TAACACACAT TAATTGCGTT GCGTCACTG CCCGCTTTC AGTCGGGAAA CCTGTCGTGC 240
CAGCTGCATT AATGAATCGG CCAACGCGC GGGAGAGGCG GTTTGCGTAT TGGGCGCTCT 300
5 TCCGCTTCCT CGTCACTGA CTCGCTGCGC TCGGTGTTT GGCTGCGGCG AGCGGTATCA 360
GCTCACTCAA AGGCGGTAAT ACGGTTATCC ACAGAATCAG GGGATAACGC AGGAAAGAAC 420
ATGTGAGCAA AAGGCCAGCA AAAGGCCAGG AACCCTAAAA AGGCCGCGTT GCTGGCGTTT 480
TTCCATAGGC TCCGCCCCC TGACGAGCAT CACAAAAATC GACGCTCAAG TCAGAGGTGG 540
CGAAACCCGA CAGGACTATA AAGATACCAG GCGTTTCCCC CTGGAAGCTC CCTCGTGCGC 600
10 TCTCCTGTTT CGACCTGCC GCTTACCGA TACCTGTCCG CTTTCTCCC TTCGGGAAGC 660
GTGGCGCTTT CTCATAGCTC ACGCTGTAGG TATCTCAGTT CCGTGTAGGT CGTTCGCTCC 720
AAGCTGGGCT GTGTGCACGA ACCCCCCGTT CAGCCCGACC GCTGCGCCTT ATCCGGTAAC 780
TATCGTCTTG AGTCCAACCC GGTAAAGACAC GACTTATCGC CACTGGCAGC AGCCACTGGT 840
AACAGGATTA GCAGAGCGAG GTATGTAGGC GGTGCTACAG AGTTCTTGAA GTGGTGGCCT 900
15 AACTACGGCT AACTAGAAG GACAGTATT GGTATCTGCG CTCTGCTGAA GCCAGTTACC 960
TTCGAAAAA GAGTTGGTAG CTCTTGATCC GGCAACAAA CCACCGCTGG TAGCGGTGGT 1020
TTTTTTGTTT GCAAGCAGCA GATTACGCG AGAAAAAAG GATCTCAAGA AGATCCTTTG 1080
ATCTTTTCTA CGGGGTCTGA CGCTCAGTG AACGAAAACT CACGTAAAG GATTTTGGT 1140
ATGAGATTAT CAAAAAGGAT CTTCACCTAG ATCCTTTTAA ATTAAAAATG AAGTTTAAA 1200
20 TCAATCTAAA GTATATATGA GTAAACTTGG TCTGACAGT ACCAATGCTT AATCAGTGAG 1260
GCACCTATCT CAGCGATCTG TCTATTTCTG TCATCCATAG TTGCTGACT CCCCCTCGTG 1320
TAGATAACTA CGATACGGA GGGCTTACCA TCTGGCCCCA GTGCTGCAAT GATACCGCGA 1380
GACCCACGCT CACCGGCTCC AGATTATCA GCAATAAACC AGCCAGCCCG AAGGGCCGAG 1440
CGCAGAAGTG GTCCTGCAAC TTTATCCGCC TCCATCCAGT CTATTAATTG TTGCCGGGAA 1500
25 GCTAGAGTAA GTAGTTCGCC AGTTAATAGT TTGCGCAACG TTGTTGCCAT TGCTACAGGC 1560
ATCGTGGTGT CACGCTCGTC GTTTGGTATG GCTTCATTCA GTCGCGTTC CCAACGATCA 1620
AGGCGAGTTA CATGATCCCC CATGTTGTGC AAAAAAGCGG TTAGCTCCTT CCGTCTCCG 1680
ATCGTTGTCA GAAGTAAGTT GGCCGCAAGT TTATCACTCA TGGTTATGGC AGCACTGCAT 1740
AATTCTCTTA CTGTCATGCC ATCCGTAAGA TGCTTTCTG TGACTGGTGA GTACTCAACC 1800
30 AAGTCATTCT GAGAATAGTG TATGCGGCGA CCGAGTTGCT CTTGCCCGGC GTCAATACGG 1860
GATAATACCG CGCCACATAG CAGAACTTTA AAAGTGCTCA TCATTGGAAA ACGTCTTCG 1920
GGGCGAAAAAC TCTCAAGGAT CTTACCGCTG TTGAGATCCA GTTCGATGTA ACCCACTCGT 1980
GCACCCAACAT GATCTTCAGC ATCTTTTACT TTCACCAGCG TTTCTGGGTG AGCAAAAACA 2040
GGAAGGCAA ATGCCGCAA AAAGGGAATA AGGCGACAC GGAAATGTTG AATACTCATA 2100
35 CTCTTCCTTT TTCAATATTA TTGAAGCATT TATCAGGGT ATTGTCTCAT GAGCGGATAC 2160
ATATTTGAAT GTATTTAGAA AAATAACAA ATAGGGGTTT CGCGCACATT TCCCGGAAA 2220
GTGCCACCTA AATTGTAAGC GTTAATATTT TGTTAAAATT CGCGTTAAAT TTTTGTTAA 2280
TCAGTCTATT TTTTAACCAA TAGGCCGAAA TCGGCAAAAT CCCTTATAAA TCAAAAGAAT 2340
AGACCGAGAT AGGGTTGAGT GTTGTTCAG TTTGGAACAA GAGTCCACTA TTAAGAAGC 2400
40 TGGACTCCAA CGTCAAAGGG CGAAAAACCG TCTATCAGGG CGATGGCCCA CTACGTGAAC 2460
CATCACCTA ATCAAGTTT TTGGGGTCTG GGTGCCGTAA AGCACTAAAT CGGAACCCTA 2520
AAGGGAGCCC CCGATTTAGA GCTTGACGGG GAAAGCCGGC GAACGTGGCG AGAAAGGAAG 2580
GGAAGAAAGC GAAAGGAGCG GCGCTAGGG CGCTGGCAAG TGTAAGCGTC ACGCTGCGCG 2640
TAACCACCAC ACCCGCCGCG CTTAATGCGC CGCTACAGGG CGCGTCCCAT TCGCCATTCA 2700
45 GGCTGCGCAA CTGTGTTGGA GGGCGATCGG TGCGGGCCTC TTCGCTATTA CGCCAGCTGG 2760
CGAAAGGGGG ATGTGCTGCA AGGCGATTAA GTTGGGTAAAC GCCAGGGTTT TCCAGTCCAC 2820
GACGTTGTAA AACGACGGCC AGTGAGCGCG CGTAATACGA CTCACTATAG GGCGAATTGG 2880
AGCTCCACCG CCGTGGCGGC CGCTCTAGAA CTAGTGATC CGTCGACTAG AGGGCCCCGAC 2940
GTCGAACCTA GGCATAAGG GATGTGAGGC CAGCATCACC GTTGACAGAAA TTGACACAAG 3000
50 CATCACCACA ATTTTCCAAA TAGAGTTTCA TTTCTTCGTC GTCAGCAGCT GCGTTGACCA 3060
TGTAATCACA CATGGAAGCC CTACACCCCA AGTTGCAATA CTTGACGGTG TCTGGTTCAT 3120
CTGAGTTGGA CACAAGGGCC AATTGGGGA AGCCTGTAGG GCATTTTCCG CTACTTGTGA 3180
GTTTACACCT ACAGACGCT GCGCATAACT TCTGAGCACC ACGGACGCGG CAAAGGTTGT 3240
AGCAGTTTCT TCCTAGGGT CTCTGCAGC AACTCTTGCC TTCTACTTGC ACCTGTTTCA 3300
55 GAACCAACCC CAGTATAAGT AAACACACCA TCACACCCTT GAGGCCCTTG CTGGTGGCCA 3360
TGG 3363

(2) INFORMATION FOR SEQ ID NO:2:

- 60 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 3365 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 65 (ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCGACCTCGA GGGGGGGCCC GGTACCCAGC TTTTGTTCCT TTAGTGAGG GTTAATTGCG 60
5 CGTTGGCGT AATCATGGTC ATAGCTGTTT CCTGTGTGAA ATTGTTATCC GCTCACAATT 120
CCACACAACA TACGAGCCGG AAGCATAAAG TGTAAGCCT GGGGTGCCTA ATGAGTGAGC 180
TAACTCACAT TAATTGCGTT GCGCTCACTG CCCGCTTCC AGTCGGGAAA CCTGTCGTGC 240
CAGCTGCATT AATGAATCGG CCAACGCGCG GGGAGAGGCG GTTTGCGTAT TGGGCGCTCT 300
TCCGTTCTCT CGCTCACTGA CTCGTTGCGC TCGGTCTGTC GGCTGCGGCG AGCGGTATCA 360
10 GCTCACTCAA AGGCGGTAAT ACGGTTATCC ACAGAATCAG GGGATAACGC AGGAAAGAAC 420
ATGTGAGCAA AAGGCCAGCA AAAGGCCAGG AACCGTAAAA AGGCCGCGTT GCTGGCGTTT 480
TTCCATAGGC TCCGCCCCC TGACGAGCAT CACAAAAATC GACGCTCAAG TCAGAGGTGG 540
CGAAACCCGA CAGGACTATA AAGATACCAG GCGTTTCCCC CTGGAAGCTC CCTCGTGCGC 600
TCTCTGTTT CGACCCTGCC GCTTACCGGA TACCTGTCCG CCTTCTCCC TTCGGGAAGC 660
15 GTGGCGCTTT CTCATAGCTC ACGCTGTAGG TATCTCAGTT CGGTGTAGGT CGTTCGCTCC 720
AAGCTGGGT GTGTGCACGA ACCCCCGTT CAGCCCGACC GCTGCGCCTT ATCCGGTAAC 780
TATCGTCTTG AGTCCAACCC GGTAAGACAC GACTTATCGC CACTGGCAGC AGCCACTGGT 840
AACAGGATTA GCAGAGCGAG GTATGTAGGC GGTGCTACAG AGTTCTTGAA GTGGTGGCCT 900
AACTACGGCT AACTAGAAG GACAGTATTT GGTATCTGCG CTCTGCTGAA GCCAGTTACC 960
20 TTCGAAAAA GAGTTGGTAG CTCTTGATCC GGCAACAAA CCACCGCTGG TAGCGGTGGT 1020
TTTTTTGTTT GCAAGCAGCA GATTACGCGC AGAAAAAAG GATCTCAAGA AGATCCTTTG 1080
ATCTTTCTA CGGGTCTGA CGCTCAGTGG AACGAAACT CACGTTAAGG GATTTTGGTC 1140
ATGAGATTAT CAAAAAGGAT CTTCACTAG ATCCTTTTAA ATTAAAAATG AAGTTTAA 1200
TCAATCTAAA GTATATATGA GTAAACTTGG TCTGACAGTT ACCAATGCTT AATCAGTGAG 1260
25 GCACCTATCT CAGCGATCTG TCTATTTCTG TCATCCATAG TTGCCTGACT CCCCCTGCTG 1320
TAGATAACTA CGATACGGA GGGCTTACCA TCTGGCCCCA GTGCTGCAAT GATACCGCGA 1380
GACCCACGCT CACCGGCTCC AGATTTATCA GCAATAAACC AGCCAGCCGG AAGGGCCGAG 1440
CGCAGAAGTG GTCCTGCAAC TTTATCCGCC TCCATCCAGT CTATTAATTG TTGCCGGGAA 1500
GCTAGAGTAA GTAGTTCGCC AGTTAATAGT TTGCGCAACG TTGTTGCCAT TGCTACAGGC 1560
30 ATCGTGGTGT CACGCTCGTC GTTTGGTATG GCTTCATTCA GCTCCGGTTC CCAACGATCA 1620
AGGCGAGTTA CATGATCCCC CATGTTGTGC AAAAAAGCGG TTAGCTCCTT CGGTCTCTCC 1680
ATCGTTGTCA GAAGTAAGTT GGCCGCACTG TTATCACTCA TGGTTATGGC AGCACTGCAT 1740
AATTCTCTTA CTGTCATGCC ATCCGTAAGA TGCTTTTCTG TGAAGGTGA GACTCAACC 1800
AAGTCATTCT GAGAATAGTG TATGCGGCGA CCGAGTTGCT CTTGCCCGGC GTCAATACCG 1860
35 GATAATACCG CGCCACATAG CAGAATTTA AAAGTGCTCA TCATTGGAAA ACGTTCTTCG 1920
GGGCGAAAAC TCTCAAGGAT CTTACCGCTG TTGAGATCCA GTTCGATGTA ACCCACTCGT 1980
GCACCAACT GATCTTCAGC ATCTTTTACT TTCACCAGCG TTTCTGGGTG AGCAAAAAACA 2040
GGAAGGCAAA ATGCCGCAAA AAAGGGAATA AGGGCGACAC GGAAATGTTG AATACTCATA 2100
CTCTTCTTTT TTCAATATTA TTGAAGCATT TATCAGGGTT ATTGTCTCAT GAGCGGATAC 2160
40 ATATTGAAT GTATTAGAA AAATAACAA ATAGGGGTTT CGCGCACATT TCCCCGAAAA 2220
GTGCCACCTA AATTGTAAGC GTTAATATTT TGTTAAAAAT CGCGTTAAAT TTTTGTAA 2280
TCAGTCAATT TTTAACCAA TAGGCCGAAA TCGGCAAAAT CCCTTATAAA TCAAAGAAT 2340
AGACCGAGAT AGGGTTGAGT GTTGTTCAG TTTGGAACAA GAGTCCACTA TTAAGAAGC 2400
TGGACTCCAA CGTCAAAGGG CGAAAAACCG TCTATCAGGG CGATGGCCCA CTACGTGAAC 2460
45 CATCACCTA ATCAAGTTT TTGGGTCGA GGTGCCGTAA AGCACTAAAT CGGAACCTA 2520
AAGGGAGCCC CGATTTAGA GCTTGACGGG GAAAGCCGGC GAACGTGGCG AGAAAGGAAG 2580
GGAAGAAAGC GAAAGGAGCG GGCCTAGGG CGCTGGCAAG TGTAAGCGGTC ACGTGCAGC 2640
TAACCAACC ACCTGCGCG CTTAATGCGC CGTACAGGG CGCGTCCAT TCGCCATTCA 2700
GGCTGCGCAA CTGTTGGGAA GGGCGATCGG TGCGGGCTC TTCGCTATTA CGCCAGCTGG 2760
50 CGAAAGGGGG ATGTGCTGCA AGGCGATTAA GTTGGGTAAC GCCAGGGTTT TCCAGTCAC 2820
GACGTTGTAA AACGACGGCC AGTGAGCGCG CGTAATACGA CTCATATAG GGCGAATTGG 2880
AGCTCCACCG CGGTGGCGGC CGCTCTAGAA CTAGTGATC CGTCGACTAG AGGGCCCGAC 2940
GTCGAACCTA GGCATAAGG GATGTGAGGC CAGCATCACC GTTGCAGAAA TTGACACAAG 3000
CATCAACCA ATTTTCCAA TAGAGTTTCA TTTCTTCGTC GTCAGCAGCT GCGTTGACCA 3060
55 TGTAAGTACA CATGGAAGCC CTACACCCCA AGTTGCAATA CTTGACGGTG TCTGGTTTCA 3120
CTGAGTTGGA CACAAGGGCC AATTGTTGGA AGCCTTTCGG GCATTTTCCG CTACTAGTCA 3180
GCTTACACTT GCAGACGCT GCGCAAAGCT TCTTGGCGCC TTTGACTTTG CAAAGGTTGT 3240
AGCACTTCT TCCAGGGTA CTCTGCAAG AACTCTTGC TTTACTTGC ACCTGTTTCA 3300
60 GAACCAACCC CAGTATAAGT AAACACACCA TCACACCTT GAGGCCCTTG CTGGTGGCCA 3360
TGGTG 3365

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

65

(A) LENGTH: 5360 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTAAATTGTA AGCGTTAATA TTTTGTIAAA ATTCGCGTTA AATTTTTGTT AAATCAGCTC 60
ATTTTTTAAC CAATAGGCCG AAATCGGCAA AATCCCTTAT AAATCAAAAG AATAGACCGA 120
10 GATAGGGTTG AGTGTGTTT CAGTTTGAA CAAGAGTCCA CTATTAAAGA ACGTGGACTC 180
CAACGTCAAA GGGCGAAAAA CCGTCTATCA GGGCGATGGC CCACTACGTG AACCATCACC 240
CTAATCAAGT TTTTGGGGT CGAGGTGCCG TAAAGCACTA AATCGGAACC CTAAGGGAG 300
CCCCCGATT AGAGCTTGAC GGGGAAAGCC GCGAACGTG GCGAGAAAGG AAGGGAAGAA 360
AGCGAAAGGA GCGGGCGCTA GGGCGCTGGC AAGTGTAGCG GTCACGCTGC GCGTAACCAC 420
15 CACACCCGCC GCGCTTAATG CGCCGCTACA GGGCGCGTCC CATTCGCCAT TCAGGCTGCG 480
CAACTGTTGG GAAGGGCGAT CGGTGCGGGC CTCTTCGCTA TTACGCCAGC TGGCGAAAGG 540
GGGATGTGCT GCAAGGCGAT TAAGTTGGGT AACGCCAGGG TTTTCCAGT CAGGACGTTG 600
TAAACGACG GCCAGTGAGC GCGCGTAATA CGACTCACTA TAGGGCGAAT TGGAGCTCCA 660
CCGCGGTGGC GGCCGCTCTA GATTATATAA TTTATAAGCT AAACAACCCG GCCCTAAAGC 720
20 ACTATCGTAT CACCTATCTA AATAAGTCAC GGGAGTTTCG AACGTCCACT TCGTCGCACG 780
GAATTGCATG TTTCTTGTTG GAAGCATATT CACGCAATCT CCACACATAA AGGTTTATGT 840
ATAAACTTAC ATTTAGCTCA GTTTAATTAC AGTCTTATTT GGATGCATAT GTATGGTTCT 900
CAATCCATAT AAGTTAGAGT AAAAAATAAG TTTAAATTTT ATCTTAATTC ACTCCAACAT 960
ATATGGATCT ACAATACTCA TGTGCATCCA AACAACTAC TTATATTGAG GTGAATTTGG 1020
25 TAGAAATTAA ACTAATTAC AACTAAGCC AATCTTTACT ATATTAAAGC ACCAGTTTCA 1080
ACGATCGTCC CGCGTCAATA TTATTAATAA ACTCCTACAT TTCTTTATAA TCAACCCGCA 1140
CTCTTATAAT CTCTTCTCTA CTAATAAT AAGAGAGTTT ATGTACAAAA TAAGGTGAAA 1200
TTATCTATAA GTGTTCTGGA TATTGGTTGT TGGTCCCAT ATTCACACAA CCTAATCAAT 1260
AGAAAACATA TGTTTTATTA AAACAAAATT TATCATATAT CATATATATA TATATATCAT 1320
30 ATATATATAT AAACCGTAGC AATGCACGGG CATATACTA GTGCAACTTA ATACATGTGT 1380
GTATTAAGAT GAATAAGAGG GTATCCAAAT AAAAACTTG TTGCTTACGT ATGGATCGAA 1440
AGGGGTGGA AACGATTAAC CGATTAAATC TCTTCCTAGT CAAAATTGAA TAGAAGGAGA 1500
TTTAATATAT CCCAATCCCC TTCGATCATC CAGGTGCAAC CGTATAAGTC CTAAGTGGT 1560
GAGGAACACG AAAGAACCAT GCATTGGCAT GTAAAGCTCC AAGAATTTGT TGTATCCTTA 1620
35 ACAATCACA GAACATCAAC AAAAATTGCA CGTCAAGGGT ATTGGGTAAG AAACAATCAA 1680
ACAAATCCTC TCTGTGCA AAGAAACAGC GTGAGTCATG CCGAGATCAT ATCATCTGA 1740
TATACATGCT TACAGCTCAC AAGACATTAC AAACACTCA TATTGCATTA CAAAGATCGT 1800
TTCATGAAAA ATAAATAGG CCGACAGGA CAAAAATCCT TGACGTGTAA AGTAAATTTA 1860
CAACAAAAAA AAAGCCATAT GTCAAGCTAA ATCTAATTCG TTTTACGTAG ATCAACAACC 1920
40 TGTAAGAGG AACAAAACTG AGCCACGCAG AAGTACAGAA TGATTCCAGA TGAACCATCG 1980
ACGTGCTACG TAAAGAGAGT GACGAGTCAT ATACATTTGG CAAGAAACCA TGAAGCTGCC 2040
TACAGCCGTC TCGGTGGCAT AAGAACACAA GAAATTGTGT TAATTAATCA AAGCTATAAA 2100
TAACGCTCGC ATGCCTGTGC ACTTCTCCAT CACCACCACT GGGTCTTCAG ACCATTAGT 2160
TTATCTACTC CAGAGCGCAG AAGAACCCGA TCGACACCAT GGCCACCAGC AAGGGCCTCA 2220
45 AGGGTGTGAT GGTGTGTTA CTTACTAGG GGTGTGTTCT CGAACAGGTG CAAGTAGAAG 2280
GCAAGAGTTG CTGCAAGAGT ACCCTGGGAA GGAAGTGCTA CAACCTTTGC AAAGTCAAAG 2340
GCGCCAAGAA GCTTTGCGCA GCGCTCTGCA AGTGTAAGCT GACTAGTAGC GGAAATGCC 2400
CGAAAGGCTT CCCCAAATTG GCCCTTGTGT CCAACTCAGA TGAACCAGAC ACCGTCAAGT 2460
ATTGCAACTT GGGGTGTAGG GCTTCCATGT GTGACTACAT GGTCAACGCA GCTGCTGACG 2520
50 ACGAAGAAAT GAAACTCTAT TTGGAAAATT GTGGTGATGC TTGTGTCAAT TTCTGCAACG 2580
GTGATGCTGG CCTCACATCC CTTAGTGCCT AAGTTCGACG TCGGGCCCTC TAGTCGACGG 2640
ATCCCCGGCG GTGTCCCCCA CTGAAGAAAC TATGTGCTGT AGTATAGCCG CTGCCCCGCTG 2700
GCTAGCTAGC TAGTTGAGTC ATTAGCGGC GATGATTGAG TAATAATGTG TCACGCATCA 2760
CCATGCATGG GTGGCAGTGT CAGTGTGAGC AATGACCTGA ATGAACAATT GAAATGAAAA 2820
55 GAAAAAGTA TTGTTCCAAA TTAACGTTTT TAACCTTTTA ATAGGTTTAT ACAATAATTG 2880
ATATATGTTT TCTGTATATG TCTAATTTGT TATCATCCAT TTAGATATAG ACAAAAAAAA 2940
ATCTAAGAAC TAAACAAAAT GCTAATTTGA AATGAAGGGA GTATATATTG GGATAATGTC 3000
GATGAGATCC CTCGTAATAT CACCGACATC ACACGTGTCC AGTTAATGTA TCAGTGATAC 3060
GTGTATTAC ATTTGTTGCG CGTAGGCGTA CCCAACAATT TTGATCGACT ATCAGAAAGT 3120
60 CAACGGAAGC GAGTCGACCT CGAGGGGGGG CCCGGTACCC AGCTTTTGTT CCCTTTAGTG 3180
AGGGTTAATT GCGCGCTTGG CGTAATCATG GTCATAGCTG TTTCTGTGT GAAATTTGTTA 3240
TCCGCTCACA ATTCCACACA ACATACGAGC CGGAAGCATA AAGTGTAAGG CTTGGGGTGC 3300
CTAATGAGTG AGCTAATCA CATTAAATTG GTTGCGCTCA CTGCCCCGTT TCCAGTCGGG 3360
AAACCTGTCG TGCCAGCTGC ATTAATGAAT CGGCCAACGC GCGGGGAGAG GCGGTTTTCG 3420
65 TATTGGGCGC TCTCCGCTT CCTCGCTCAC TGACTCGCTG CGCTCGGTG TCCGGCTGCG 3480
GCGAGCGGTA TCAGCTCACT CAAAGGCGGT AATACGGTTA TCCACAGAAT CAGGGGATAA 3540

CGCAGGAAAG AACATGTGAG CAAAAGGCCA GCAAAGGCC AGGAACCGTA AAAAGGCCGC 3600
GTTGCTGGCG TTTTCCATA GGCTCCGCC CCCTGACGAG CATCACAAAA ATCGACGCTC 3660
AAGTCAGAGG TGGCGAAACC CGACAGGACT ATAAAGATAC CAGGCGTTTC CCCCTGGAAG 3720
CTCCCTCGTG CGCTCTCCTG TTCCGACCCT GCCGCTTACC GGATACCTGT CCGCCTTTCT 3780
5 CCCTTCGGGA AGCGTGGCGC TTTCTCATAG CTCACGCTGT AGGTATCTCA GTTCGGTGTA 3840
GGTCGTTGCG TCCAAGCTGG GCTGTGTGCA CGAACCCCC GTTCAGCCCG ACCGCTGCGC 3900
CTTATCCGGT AACTATCGTC TTGAGTCCAA CCCGGTAAGA CACGACTTAT CGCCACTGGC 3960
AGCAGCCACT GGTAACAGGA TTAGCAGAGC GAGGTATGTA GGCGGTGCTA CAGAGTTCTT 4020
GAAGTGGTGG CCTAACTACG GCTACACTAG AAGGACAGTA TTTGGTATCT GCGCTCTGCT 4080
10 GAAGCCAGTT ACCTTCGGAA AAAGAGTTGG TAGCTCTTGA TCCGGCAAAC AAACCACCGC 4140
TGGTAGCGGT GGTTTTTTTT TTTGCAAGCA GCAGATTACG CGCAGAAAAA AAGGATCTCA 4200
AGAAGATCCT TTGATCTTTT CTACGGGGTC TGACGCTCAG TGAACGAAA ACTCACGTTA 4260
AGGGATTTTG GTCATGAGAT TATCAAAAAG GATCTTCACC TAGATCCTTT TAAATTAATA 4320
ATGAAGCTT AAATCAATCT AAAGTATATA TGAGTAAACT TGGTCTGACA GTTACCAATG 4380
15 CTTAATCAGT GAGGCACCTA TCTCAGCGAT CTGTCTATTT CGTTCATCCA TAGTTGCCTG 4440
ACTCCCCGTC GTGTAGATAA CTACGATACG GGAGGGCTTA CCATCTGGCC CCAGTGCTGC 4500
AATGATACCG CGAGACCCAC GCTCACCGGC TCCAGATTTA TCAGCAATAA ACCAGCCAGC 4560
CGGAAGGGCC GAGCGCAGAA GTGGTCTGCA AACTTTATCC GCCTCCATCC AGTCTATTAA 4620
TTGTTGCCGG GAAGCTAGAG TAAGTAGTTC GCCAGTTAAT AGTTTGCACA ACGTTGTTGC 4680
20 CATTGCTACA GGCATCGTGG TGTACGCTC GTCGTTTGGT ATGGCTTCAT TCAGTCCGG 4740
TTCCCAACGA TCAAGGCGAG TTACATGATC CCCCATGTTG TGCAAAAAAG CGGTTAGCTC 4800
CTTCGGTCTT CCGATCGTTG TCAGAAGTAA GTTGGCCGCA GTGTTATCAC TCATGGTTAT 4860
GGCAGCACTG CATAATTCTC TTAGTGTCAT GCCATCCGTA AGATGCTTTT CTGTGACTGG 4920
TGAGTACTCA ACCAAGTCAT TCTGAGAATA GTGTATGCGG CGACCGAGTT GCTCTTGCCC 4980
25 GCGTCAATA CGGGATAATA CCGCGCCACA TAGCAGAACT TAAAAGTGC TCATCATTGG 5040
AAAACGTTCT TCGGGGCGAA AACTCTCAAG GATCTTACCG CTGTTGAGAT CCAGTTTCAT 5100
GTAACCCACT CGTGACCCA ACTGATCTTC AGCATCTTTT ACTTTCACCA GCGTTTCTGG 5160
GTGAGCAAAA ACAGGAAGGC AAAATGCCGC AAAAAAGGGA ATAAGGGCGA CACGGAATG 5220
TTGAATACTC ATACTCTTCC TTTTCAATA TTATTGAAGC ATTTATCAGG GTTATTGTCT 5280
30 CATGAGCGGA TACATATTTG AATGTATTTA GAAAAATAAA CAAATAGGGG TTCCGCGCAC 5340
ATTTCCCCGA AAAGTGCCAC 5360

(2) INFORMATION FOR SEQ ID NO:4:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5511 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

45 TCGCGCGTTT CGGTGATGAC GGTGAAAACC TCTGACACAT GCAGCTCCCG GAGACGGTCA 60
CAGCTTGTCT GTAAGCGGAT GCCGGGAGCA GACAAGCCCG TCAGGGCGCG TCAGCGGGTG 120
TTGGCGGGTG TCGGGGCTGG CTTAACTATG CGGCATCAGA GCAGATTGTA CTGAGAGTGC 180
ACCATATGCG GTGTGAAATA CCGCACAGAT GCGTAAGGAG AAAATACCGC ATCAGGCGCC 240
ATTCGCCATT CAGGCTGCGC AACTGTTGGG AAGGGCGATC GGTGCGGGCC TCTTCGCTAT 300
50 TACGCCAGCT GCGGAAAGGG GGATGTGCTG CAAGGCGATT AAGTTGGGTA ACGCCAGGGT 360
TTTCCAGTC ACGACGTTGT AAAACGACGG CAGTGAATT CTTTATGAA TAATAATAAT 420
GCATATCTGT GCATTACTAC CTGGGATACA AGGGCTTCTC CGCCATAACA AATTGAGTTG 480
CGATGCTGAG AACGAACGGG GAAGAAAGTA AGCGCCGCC AAAAAGGCG AACATGTACG 540
TCGGCTATAG CAGGTGAAAG TTCGTGCGCC AATGAAAAGG GAACGATATG CGTTGGGTAG 600
55 TTGGGATACT TAAATTTGGA GAGTTTGTG CATACTAA TCCACTAAAG TTGTCTATCT 660
TTTAAACAGC TCTAGGCAGG ATATAAGATT TATATCTAAT CTGTTGGAGT TGCTTTTAGA 720
GTAACTTTTC TCTCTGTTT GTTTATAGCC GATTAGCACA AAATTAACT AGGTGACGAG 780
AAATAAAGAA AAACGGAGGC AGTAAAAAAT ACCCAAAAAA ATACTTGGAG ATTTTGTCT 840
CAAAATTATC TTCTAATTTT AAAAGCTACA TATTAATAAT ACTATATATT AAAAAATG 900
60 CGAGATCATT GCTTGGGATG GGCAGGGCCA ATAGCTAATT GCTAAGGATG GGCTATATTT 960
ATGTATCGTC TGAAACATGT AGGGGCTAAT AGTTAGATGA CTAATTTGCT GTGTTCGTAC 1020
GGGGTGCTGT TTGAGCCTAG CGATGAAGGG TCATAGTTTC ATACAAGAAC TCACTTTTGG 1080
TTCGTCTGCT GTGTCTGTT TCAGCGTAAC GGCATCAATG GATGCCAAAC TCCGAAGGG 1140
GACAAATGAA GAAGCGAAGA GATTATAGAA CACGCACGTG TCATTATTTA TTTATGGACT 1200
65 TGCTTCAGTA GCTTACAGCA TCGTACCGC ACGTACATAC TACAGAGCCA CACTTATTGC 1260
ACTGCCTGCC GCTTACGTAC ATAGTTAACA CGCAGAGAGG TATATACATA CACGTCCAAC 1320

GTCTCCACTC AGGCTCATGC TACGTACGCA CGTCGGTCGC GCGCCACCCT CTCGTTGCTT 1380
 CCTGCTCGTT TTGGCGAGCT AGAGGGCCCC ACGTCGAACT TAGGCACTAA GGGATGTGAG 1440
 GCCAGCATCA CCGTTGCAGA AATTGACACA AGCATACCA CAATTTTCCA AATAGCGTTT 1500
 CATTCTTCG TCGTCAGCAG CTGCGTTGAC CATGTAGTCA CACATGGAAG CCTACACCC 1560
 5 CAAGTTGCAA TACTTGACGG TGTCTGGTTC ATCTGAGTTG GACACAAGGG CCAATTTGGG 1620
 GAAGCCTTTC GGGCATTTTC CGCTACTAGT CAGCTTACAC TTGCAGACGC CTGCGCAAAG 1680
 CTTCTGGCG CTTTGACTT TGCAAAGGTT GTAGCACTTC CTTCCAGGG TACTCTTGCA 1740
 GCAACTCTTG CTTTCTACTT GCACCTGTTT GAGAACCAAC CCCAGTATAA GTAAACACAC 1800
 CATCACACCC TTGAGGCCCT TGCTGGTGGC CATGGTGTAG TGTCGACTGT GATATCCTCG 1860
 10 GGTGTGTGTT TGATCCTTGG GTTGGCTGTA TGCAGACTA AAGCGGAGGT GGCGCGATT 1920
 TATACCAGCG CCGGGCCCTG GTACGTGGCG CGGCCGCGCG GCTACGTGGA GGAAGGCTGC 1980
 GTGGCAGCAG ACACACGGGT CGCCACGTCC CGCCGTACTC TCCTTACCGT GCTTATCCGG 2040
 GCTCCGGCTC GGTGCACGCC AGGGTGTGGC CGCCTCTGAG CAGACTTTGT CGTGTTCAC 2100
 AGTGGTGTG TGTTCGGGG ACTCCGATCC GCGGCGAGCG ACCGAGCGTG TAAAAGAGTT 2160
 15 CCTACTAGT ACGTTCATTG TATCTGGACG ACGGGCAGCG GACAATTTGC TGTAAGAGAG 2220
 GGGCAGTTTT TTTTAGAAA AACAGAGAAT TCCGTTGAGC TAATTGTAAT TCAACAAATA 2280
 AGCTATTAGT TGGTTTTAGC TTAGATTAAG GAAGCTAACG ACTAATAGCT AATAATTAGT 2340
 TGGTCTATTA GTTGACTCAT TTTAAGGCCG TGTTTCAATC TCGCGAGATA AACTTTAGCA 2400
 GCTATTTTTT AGTACTTTT AGCCATTTGT AATCTAAACA GGAGAGCTAA TGGTGGTAAT 2460
 20 TGAAACTAAA CTTTAGCACT TCAATTCATA TAGCTAAAGT TTAGCAGGAA GCTAAACTTT 2520
 ATCCCGTGAG ATTGAAACGG GGCCTAAATC TCTCAGCTAT TTTTGATGCA AATTACTGTC 2580
 ACTACTGGAA TTGAGCGCTT TGCCGAGTGT CAAAGCCTGA AAAACACTCC GTAAAGACT 2640
 TGCCTAGTGT GACACTCGAC AAAGAGATCT CGACGAACAG TACATCGACA ACGGCTTCTT 2700
 TGTCGAGTAC TTTTATCGG AACTTGACA AAGTCTTGT CGAGTGAAT ACATTGAAAC 2760
 25 TCTATGATT TATGTGTAGG TCACTTAGGT TTCTACACAT AGTACGTCAC AACTTTACCG 2820
 AAACATTATC AAATTTTTAT CACAACCTCT ATATATGATA TCATGACATG TGGACAAGTT 2880
 TCATTAAATT CTGACTTTAT TTGTGTTTTA TACAATTTTT AAACAAGTAT ATAACAAGTT 2940
 CACGGTCATG TTTAGTGAGC ATGGTGCTTG AAGATTCTGG TCTGCTTCTG AAATCGGTCG 3000
 TAACTTGTGC TAGATAACAT GCATATCATT TATTTTGCAT GCACGGTTTT CCATGTTTTG 3060
 30 AGTGAATGC AGTTTAAATG TGAATTTTCC GAAGAAATTC AAATAAACGA ACTAAATCTA 3120
 ATATTTATAG AAAACATTTT TGTAATATG TAATTGTGCC AAAATGGTAC ATGTAGATCT 3180
 ACATAGTGTA GGAACATACC AAAAAAGTT TGGTTGGCAA AATAAAAAAA ATAAATATA 3240
 CTTTATCGAG TGTCCAAGGA TGGCACTCGG CAAGCTTGGC GTAATCATGG TCATAGCTGT 3300
 TTCTGTGTG AAATTGTTAT CCGCTCACA TTCCACACAA CATACGAGCC GGAAGCATAA 3360
 35 AGTGTAAAGC CTGGGGTGCC TAATGAGTGA GCTAACTCAC ATTAATTGCG TTGCGCTCAC 3420
 TGCCCGCTT CCAGTCGGGA AACCTGTCTG GCCAGCTGCA TTAATGAATC GGCCAACGCG 3480
 CGGGGAGAGG CGGTTTGCCT ATTGGGCGCT CTTCCGCTC CTCGCTCACT GACTCGCTGC 3540
 GCTCGGTCTG TCGGCTGCGG CGAGCGGTAT CAGCTCACTC AAAGGCGGTA ATACGGTTAT 3600
 CCACAGATA AGGGGATAAC GCAGGAAAGA ACATGTGAGC AAAAGGCCAG CAAAAGGCCA 3660
 40 GGAACCGTAA AAAGGCCGCG TTGTGGCGT TTTTCCATAG GCTCCGCCCC CCGACGAGC 3720
 ATCACAATAA TCGACGCTCA AGTCAGAGGT GGCGAAACCC GACAGGACTA TAAAGATACC 3780
 AGGCGTTTCC CCCTGGAAGC TCCCTCGTGC GCTCTCTGT TCCGACCCTG CCGCTTACCG 3840
 GATACCTGTC CGCCTTTCTC CTTTCGGGAA GCGTGGCGCT TTCTCAATGC TCACGCTGTA 3900
 GGTATCTCAG TTCGGTGTAG GTCGTTTCGT CCAAGCTGGG CTGTGTGCAC GAACCCCCCG 3960
 45 TTCAGCCGA CCGCTGCGCC TTATCCGGTA ACTACTGCT TGAGTCCAAC CCGGTAAGAC 4020
 ACCGATTATC GCCACTGGCA CGACGACTG GTAACAGGAT TAGCAGAGCG AGGTATGTAG 4080
 GCGGTGCTAC AGAGTTCTTG AAGTGGTGGC CTAACACGG CTACACTAGA AGGACAGTAT 4140
 TTGGTATCTG CGCTCTGCTG AAGCCAGTTA CTTTCGGAAA AAGAGTTGGT AGCTCTTGAT 4200
 CCGGCAAAAC AACCACCGCT GGTAGCGGTG GTTTTTTGT TTGCAAGCAG CAGATTACGC 4260
 50 GCAGAAAAAA AGGATCTCAA GAAGATCCTT TGATCTTTT TACGGGGTCT GACGCTCAGT 4320
 GGAACGAAAA CTCACGTAA GGGATTTTGG TCATGAGATT ATCAAAAAGG ATCTTCACCT 4380
 AGATCTTTT AAATTAATAA TGAAGTTTTA AATCAATCTA AAGTATATAT GAGTAAACTT 4440
 GGTCTGACAG TTACCAATGC TTAATCAGTG AGGCACCTAT CTCAGCGATC TGTCTATTTC 4500
 GTTCATCCAT AGTTGCCTGA CTCCCCGTCG TGTAAGTAAAC TACGATACGG GAGGGCTTAC 4560
 55 CATCTGGCCC CAGTGCTGCA ATGATACCGC GAGACCCACG CTCACCGGCT CCAGATTTAT 4620
 CAGCAATAAA CCAGCCAGCC GGAAGGGCCG AGCGCAGAAG TGGTCCTGCA ACTTTATCCG 4680
 CCTCCATCCA GTCTATTAAT TGTTGCCGGG AAGCTAGAGT AAGTAGTTTC CCAGTTAATA 4740
 GTTTGCGCAA CGTTGTTGCC ATTGCTACAG GCATCGTGGT GTCACGCTCG TCGTTTGGTA 4800
 TGGCTTCATT CAGCTCCGGT TCCCAACGAT CAAGGCGAGT TACATGATCC CCCATGTTGT 4860
 60 GCAAAAAAGC GGTTAGCTCC TTCGGTCTC CGATCGTTGT CAGAAGTAAG TTGGCCGCAG 4920
 TGTTATCACT CATGGTTATG GCAGCACTGC ATAATCTCT TACTGTCATG CCATCCGTAA 4980
 GATGCTTTT TGTGACTGGT GAGTACTCAA CCAAGTCATT CTGAGAATAG TGATGCGGC 5040
 GACCGAGTTG CTCTTGCCCG GCGTCAATAC GGGATAATAC CGCGCCACAT AGCAGAAGTT 5100
 TAAAGATGCT CATCATTTGA AAACCTTCTT CGGGCGGAAA ACTCTCAAGG ATCTTACCGC 5160
 65 TGTTGAGATC GATTCGATG TAACCCACTC GTGCACCCAA CTGATCTTCA GCATCTTTTA 5220
 CTTTACCAG CGTTTCTGGG TGAGCAAAAA CAGGAAGGCA AAATGCCGCA AAAAAGGGAA 5280

TAAGGGCGAC ACGGAAATGT TGAATACTCA TACTCTTCCT TTTTCAATAT TATTGAAGCA 5340
TTTATCAGGG TTATTGTCTC ATGAGCGGAT ACATATTTGA ATGTATTTAG AAAAATAAAC 5400
AAATAGGGGT TCCGCGCACA TTTCCCCGAA AAGTGCCACC TGACGTCTAA GAAACCATTA_5460
TTATCATGAC ATTAACCTAT AAAAATAGGC GTATCACGAG GCCCTTTCGT C 5511

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5115 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GTTGGGAGCT CTCCCATATG GTCGACCTGC AGGCGGCCGC TCTAGAACTA GTGGATCCCC 60
CCCTCGAGGT CGACGGTATC GATAAGCTTG ATATCTTACA AGGCCAGCC CAGCGACCTA 120
TTACACAGCC CGCTCGGGCC CGCGACGTGC GGACACATCT TCTTCCCCCT TTTGGTGAAG 180
CTCTGCTCGC AGCTGTCCGG CTCCTTGGAC GTTCGTGTGG CAGATTCATC TGTGTCTCG 240
TCTCCTGTGC TTCTGTGGTA GCTTGTGTAG TGGAGCTGAC ATGGTCTGAG CAGGCTTAAA 300
ATTTGCTCGT AGACGAGGAG TACCAGCACA GCACGTTGCG GATTTCTCTG CCTGTGAAGT 360
GCAACGTCTA GGATTGTCAC ACGCCTTGGT CGCGTCGCGT CGCGTCGCGT CGATGCGGTG 420
GTGAGCAGAG CAGCAACAGC TGGGCGGCC AACGTTGGCT TCCGTGTCTT CGTCGTACGT 480
ACGCGCGCGC CGGGGACACG CAGCAGAGAG CGGAGAGCGA GCCGTGCACG GGGAGGTGGT 540
GTGGAAGTGG AGCCGCGCGC CCGGCCGCC GCGCCCGGTG GGCAACCCAA AAGTACCCAC 600
GACAGCGAA GCGGCCAAAAG CGATCCAAGC TCCGGAACGC AACAGCATGC GTCGCGTCGG 660
AGAGCCAGCC ACAAGCAGCC GAGAACCGAA CCGGTGGGCG ACGCGTCATG GGACGGACGC 720
GGGCGACGCT TCCAAACGGG CCACGTACGC CGGCGTGTGC GTGCGTGCAG ACGACAAGCC 780
AAGGCGAGGC AGCCCCCGAT CGGGAAGCG TTTTGGGCGC GAGCGCTGGC GTGCGGGTCA 840
GTCGCTGGTG CGCAGTGCCG GGGGGAACGG GTATCGTGGG GGGCGCGGGC GGAGGAGAGC 900
GTGGCGAGGG CCGAGAGCAG CGCGCGGCCG GGTACGCAA CGCGCCCCAC GTACTGCCCT 960
CCCCCTCCG CCGCGCTAGA AATACCGAGG CCTGGACCGG GGGGGGGGCC CGTCACATCC 1020
ATCCATCGAC CGATCGATCG CCACAGCCAA CACCACCCGC CGAGGCGACG CGACAGCCGC 1080
CAGGAGGAAG GAATAAACTC ACTGCCAGCC AGTGAAGGGG GAGAAGTGTA CTGCTCCGTC 1140
GACCAGTGC CGCACCGCCC GGCAGGGCTG CTCATCTCGT CGACGACCAG GTTCTGTTCC 1200
GATCCGATCC GATCCTGTCC TTGAGTTTCG TCCAGATCCT GCGCGGTATC TCGTGTTT 1260
ATGATCCAGG TTCTTCGAAC CTAATCTGT CCGTGCACAC GTCTTTTCTC TCTCTCTAC 1320
GCAGTGGATT AATCGCCATG GCCACCAGCA AGGCCCTCAA GGGTGTGATG GTGTGTTTAC 1380
TTATACTGGG GTTGGTTCTC GAACAGGTGC AAGTAGAAGG CAAGAGTTGC TGCAAGAGTA 1440
CCCTGGGAAG GAAGTGCTAC AACCTTTGCA AAGTCAAAGG CGCCAAGAAG CTTTGCGCAG 1500
GCGTCTGCAA GTGTAAGCTG ACTAGTAGCG GAAAATGCC GAAAGGCTTC CCAAATTGG 1560
CCCTGTGTG CAACTCAGAT GAACCAGACA CCGTCAAGTA TTGCAACTTG GGGTGTAGGG 1620
CTTCCATGTG TGACTACATG GTCAACGCAG CTGCTGACGA CGAAGAAATG AAACCTTATT 1680
TGGAATAATTG TGGTATGCT TGTGTCAATT TCTGCAACGG TGATGCTGGC CTCACATCCC 1740
TTAGTGCTTA AGTTCGACGT CGGGCCCTCT AGATGCGGCC CGGGTGAAGA GTTCGCCCTG 1800
CAGGGCCCCT GATCTCGCG GTGGTGCAA GATGTTGGGA CATCTTCTTA TATATGCTGT 1860
TTCGCTTATG TGATATGGAC AAGTATGTGT AGATGCTTGC TTGTGCTAGT GTAATGTAGT 1920
GTAGTGGTGG CAGTGGCAC AACCTAATAA GCGCATGAAC TAATTGCTTG CGTGTGTAGT 1980
TAAGTACCGA TCGGTAATTT TATATTGCGA GTAAATAAAT GGACCTGTAG TGGTGGAGTA 2040
AATAATCCCT GCTGTCGGT GTTCTTATCG TCAATCGTAT AGATATTATA TAGAGTACAT 2100
TTTTCTCTCT CTGAATCCTA CGTGTGTGAA ATTTCTATAT CATTACTGTA AAATTTCTGC 2160
GTTCCAAAAG AGACCATAGC CTATCTTTGG CCCTGTTTGT TTCGGCTTCT GGCAGCTTCT 2220
GGCCACCAA AGCTGCTGCG GACTGCCAAA CGCTCAGATT TTCAGTAGC TTCTATAAAA 2280
TTAGTTGGGG CAAAACCAT CCAAATCAA TATAACACA TAATCGGTTG AGTCGTTGTA 2340
ATATTAGGAA TCTGTCACTT TCTAGATCCT GAGCCCTATG AACAACCTTA TCTTTCTCCA 2400
TACGTAATCG TAATGATACT CAGATTCTCT CCACAGCCAG ATTCTCCTCA CAGCCAGATT 2460
TTCAGAAAAG CTGGTCAGAA AAAAGTTAAA CCAAACAGAC CCTTTGTGTA TGCATGGAT 2520
GGCTTTCCCC GTCAAGCTCT AAATCGGGGG CTCCTTTAG GGTTCGATT TAGAGCTTTA 2580
CGGCACCTCG ACCGCAAAAA ACTTGATTG GGTGATGGT CACGTAGTG GCCATCGCCC 2640
TGATAGACGG TTTTTCGCCC TTTGACGTTG GAGTCCACGT TCTTTAATAG TGGACTCTG 2700
TTCCAAACTG GAACAACACT CAACCTATC TCGGTCTATT CTTTGTATT ATAAGGGATT 2760
TTGCCGATT CGGCCTATT GTTAAAAAT GAGCTGATT AACAAATATT TAACGCGAAT 2820
TTTAAACAAA TATTAACGTT TACAATTCG CCTGATGCGG TATTTCTCC TTACGCATCT 2880
GTGCGGTATT TCACACCGCA TACAGGTGGC ACTTTTCGGG GAAATGTGCG CGGAACCCCT 2940

ATTTGTTTAT TTTTCTAAAT ACATTCAAAT ATGTATCCGC TCATGAGACA ATAACCCTGA 3000
 TAAATGCTTC AATAATATTG AAAAAGGAAG AGTATGAGTA TTCAACATTT CCGTGTGCGC 3060
 CTTATTCCCT TTTTTCGGC ATTTTGCCTT CCTGTTTTTG CTCACCCAGA AACGCTGGTG 3120
 5 AAAGTAAAAG ATGCTGAAGA TCAGTTGGGT GCACGAGTGG GTTACATCGA ACTGGATCTC 3180
 AACAGCGGTA AGATCCTTGA GAGTTTTTCG CCCGAAGAAC GTTTTCCAAT GATGAGCACT 3240
 TTAAAGTTC TGCTATGTCA TACACTATTA TCCCGTATTG ACGCCGGGCA AGAGCAACTC 3300
 GGTGCGCGGG CGCGGTATTC TCAGAAATGAC TTGGTTGAGT ACTCACCAGT CACAGAAAAG 3360
 CATCTTACGG ATGGCATGAC AGTAAGAGAA TTATGCAGTG CTGCCATAAC CATGATGAT 3420
 AACACTGCGG CCAACTTACT TCTGACAACG ATCGGAGGAC CGAAGGAGCT AACCGCTTTT 3480
 10 TTGCACAACA TGGGGGATCA TGTAATCTCG CTTGATCGTT GGAACCGGA GCTGAATGAA 3540
 GCCATACCAA ACGACGAGCG TGACACCACG ATGCCTGTAG CAATGCCAAC AACGTTGCGC 3600
 AAATATTAA CTGGCGAACT ACTTACTCTA GCTTCCCGGC AACAATTAAT AGACTGGATG 3660
 GAGGCGGATA AAGTTGCAGG ACCACTTCTG CGCTCGGCCC TTCCGGCTGG CTGGTTTATT 3720
 GCTGATAAAT CTGGAGCCGG TGAGCGTGGG TCTCGCGGTA TCATTGCAGC ACTGGGGCCA 3780
 15 GATGGTAAGC CCTCCCGTAT CGTAGTTATC TACACGACGG GGAGTCAGGC AACTATGGAT 3840
 GAACGAAATA GACAGATCGC TGAGATAGGT GCCTCACTGA TTAAGCATTG GTAACGTGCA 3900
 GACCAAGTTT ACTCATATAT ACTTTAGATT GATTTAAAAAC TTCATTTTTA ATTTAAAAGG 3960
 ATCTAGGTGA AGATCCTTTT TGATAATCTC ATGACCAAAA TCCCTTAACG TGAGTTTTTCG 4020
 TTCCACTGAG CGTCAGACCC CGTAGAAAAG ATCAAAGGAT CTTCTTGAGA TCCTTTTTTT 4080
 20 CTGCGCGTAA TCTGCTGCTT GCAAAACAAA AAACCACCGC TACCAGCGGT GGTGTTGTTG 4140
 CCGGATCAAG AGCTACCAAC TCTTTTTCCG AAGGTAAGT GCTTCAGCAG AGCGCAGATA 4200
 CCAAATACTG TCCTTCTAGT GTAGCCGTAG TTAGGCCACC ACTTCAAGAA CTCTGTAGCA 4260
 CCGCTACAT ACCTCGCTCT GCTAATCTG TTACCAGTGG CTGCTGCCAG TGGCGATAAG 4320
 TCGTGTCTTA CCGGTTTGA CTCAAGACGA TAGTTACCGG ATAAGGCGCA GCGGTCGGGC 4380
 25 TGAACGGGGG GTTCGTGCAC ACAGCCCAGC TTGGAGCGAA CGACCTACAC CGAAGTGAAG 4440
 TACCTACAGC GTGAGCTATG AGAAAGCGCC ACGCTTCCCG AAGGGAGAAA GGCGGACAGG 4500
 TATCCGGTAA CCGGCAGGGT CGGAACAGGA GAGCGCACGA GGGAGCTTCC AGGGGGAAAC 4560
 GCCTGGTATC TTATAGTCC TGTCGGGTTT CGCCACCTCT GACTTGAGCG TCGATTTTTG 4620
 TGATGCTCGT CAGGGGGGCG GAGCCTATCG AAAAACGCCA GCAACGCGGC CTTTTTACGG 4680
 30 TTCCTGGCCT TTTGCTGGCC TTTTGCTCAC ATGTTCTTTC CTGCGTTATC CCCTGATTCT 4740
 GTGGATAACC GTATTACCGC CTTTGAGTGA GCTGATACCG CTCGCCGAG CCGAACGACC 4800
 GAGCGCAGCG AGTCAGTGAG CGAGGAAGCG GAAGAGCGCC CAATACGCAA ACCGCCTCTC 4860
 CCGCGCGTGG GGCCGATTCA TTAATGCAGC TGGCAGCACA GGTTCCTCGA CTGGAAAGCG 4920
 GGCAGTGAGC CGAACGCCAAT TAATGTGAGT TAGCTCACTC ATTAGGCACC CCAGGCTTTA 4980
 35 CACTTTATGC TTCCGGCTCG TATGTTGTGT GGAATTGTGA GCGGATAACA ATTTACACACA 5040
 GGAAACAGCT ATGACCATGA TTACGCCAAG CTATTTAGGT GACACTATAG AATACTCAAG 5100
 CTATGCATCC AACGC 5115

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5392 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CTAAATTGTA AGCGTTAATA TTTTGTTAAA ATTCGCGTTA AATTTTTGTT AAATCAGCTC 60
 ATTTTTTAAC CAATAGGCCG AAATCGGCAA AATCCCTTAT AAATCAAAAAG AATAGACCGA 120
 GATAGGGTTG AGTGTGTTC CAGTTTGGA CAAGAGTCCA CTATTAAAGA ACGTGGACTC 180
 CAACGTCAAA GGGCGAAAAA CCGTCTATCA GGGCGATGGC CCACTACGTG AACCATCACC 240
 55 CTAATCAAGT TTTTGGGGT CGAGGTGCCG TAAAGCACTA AATCGGAACC CTAAGGGGAG 300
 CCCCCGATT AGAGCTTGAC GGGGAAAGCC GGCGAACGTG GCGAGAAAGG AAGGGAAGAA 360
 AGCGAAAGGA GCGGGCGCTA GGGCGCTGGC AAGTGTAGCG GTCACGCTGC GCGTAACCAC 420
 CACACCCGCC GCGCTTAATG CGCCGCTACA GGGCGCGTCC CATTGCCAT TCAGGCTGCG 480
 CAAGTGTGG GAAGGGCGAT CGGTGCGGGC CTCTTCGTA TTACGCCAGC TGGCGAAAGG 540
 60 GGGATGTGCT GAAAGCGAT TAAGTTGGGT AACGCCAGG TTTCCAGT CACGACGTTG 600
 TAAACGACG GCCAGTGAGC GCGCGTAATA CGACTCACTA TAGGGCGAAT TGGAGCTCCA 660
 CCGCGTGGC GGCCGCTCTA GATTATATAA TTTATAAGCT AAACAACCCG GCCCTAAAGC 720
 ACTATCGTAT CACCTATCTA AATAAGTCAC GGGAGTTTCG AACGTCCACT TCGTCGCACG 780
 GAATTGCATG TTTCTTGTG GAAGCATATT CACGCAATCT CCACACATAA AGGTTTATGT 840
 65 ATAACTTAC ATTTAGCTCA GTTTAATTAC AGTCTTATTT GGATGCATAT GTATGGTTCT 900
 CAATCCATAT AAGTTAGAGT AAAAAATAAG TTAAATTTT ATCTTAATTC ACTCCAACAT 960

ATATGGATCT ACAATACTCA TGTGCATCCA AACAACTAC TTATATTGAG GTGAATTTGG 1020
TAGAAATTAA ACTAATTAC ACTAAGCC AATCTTTACT ATATTAAAGC ACCAGTTTCA 1080
ACGATCGTCC CGCGTCAATA TTATTAATAA ACTCCTACAT TTCTTTATAA TCAACCCGCA 1140
CTCTTATAAT CTCTTCTCTA CTACTATAAT AAGAGAGTTT ATGTACAAAA TAAGGTGAAA 1200
5 TTATCTATAA GTGTTCTGGA TATTGGTTGT TGGCTCCCAT ATTCACACAA CCTAATCAAT 1260
AGAAAACATA TGTTTTATTA AAACAAAATT TATCATATAT CATATATATA TATATATCAT 1320
ATATATATAT AAACCGTAGC AATGCACGGG CATATAACTA GTGCAACTTA ATACATGTGT 1380
GTATTAAGAT GAATAAGAGG GTATCCAAAT AAAAACTTG TTGCTTACGT ATGGATCGAA 1440
AGGGGTTGGA AACGATTAAA CGATTAAATC TCTTCTAGT CAAAATTGAA TAGAAGGAGA 1500
10 TTTAATATAT CCCAATCCCC TTCGATCATC TCGTGCAAC CGTATAAGTC CTAAGTGGT 1560
GAGGAACACG AAAGAACCAT GCATTGGCAT GTAAAGCTCC AAGAATTTGT TGTATCCTTA 1620
ACAACTCACA GAACATCAAC CAAAATTGCA CGTCAAGGGT ATTGGGTAAG AAACAATCAA 1680
ACAAATCCTC TCTGTGTGCA AAGAAACACG GTGAGTCATG CCGAGATCAT ACTCATCTGA 1740
TATACATGCT TACAGCTCAC AAGACATTAC AAACAACCTCA TATTGCATTA CAAAGATCGT 1800
15 TTCATGAAAA ATAAATAGG CCGGACAGGA CAAAAATCCT TGACGTGTAA AGTAAATTTA 1860
CAACAAAAAA AAAGCCATAT GTCAAGCTAA ATCTAATTCG TTTTACGTAG ATCAACAACC 1920
TGTAAGAGG AACAAAACTG AGCCACGCAG AAGTACAGAA TGATTCCAGA TGAACCATCG 1980
ACGTGCTACG TAAAGAGAGT GACGAGTCAT ATACATTTGG CAAGAAACCA TGAAGCTGCC 2040
TACAGCCGTA TCGGTGGCAT AAGAACACAA GAAATTTGTG TAATTAATCA AAGCTATAAA 2100
20 TAACGCTCGC ATGCCTGTGC ACTTCTCCAT CACCACCACT GGGTCTTCAG ACCATTAGCT 2160
TTATCTACT CAGAGCGCAG AAGAACCCGA TCGACACCAT GACCAAGTTC ACAATCCTCC 2220
TCATCTCTCT TCTCTTCTGC ATCGCCACA CTTGCAGCGC CTCCAAATGG CAGCACCAG 2280
AAGATAGCTG CCGCAAGCAG CTTAAGGGGG TGAACCTCAC GCCCTGCGAG AAGCACATCA 2340
TGGAGAAGAT CCAAGGCCGC GGCGATGACG ATGATGATGA TGACGACGAC AATCACATTC 2400
25 TCAGGACCAT GCGGGGGAAG AATCACTACA TACGGAAGAA GGAAGGAAAA GACGAAGACG 2460
AAGAAGAAGA AGGACACATG CAGAAGTGCT GCGCTTTGCA CTGGCATTGT GGGCTCTTAA 2520
GCTCGTCTAT TCTGTGCTG CAGAAGATAA TGGAGAACCA GAGCGAGGAA CTGGAGGAGA 2580
AGGAGAAGAA GAAATGGAG AAGGAGCTTA TGAACCTTGGC TACTATGTGC AGGTTTGGGC 2640
CCATGATCGG GTGCGACTTG TCTCCGATG ACTAAGTTGA TCCCCGGCGG TGTCCTCCAC 2700
30 TGAAGAACT ATGTGCTGTA GTATAGCCG TGGCTAGCTA GCTAGTTGAG TCATTTAGCG 2760
GCGATGATTG AGTAATAATG TGTCACGCAT CACCATGCAT GGGTGGCAGT CTCAGTGTGA 2820
GCAATGACCT GAATGAACAA TTGAAATGAA AAGAAAAAAG TATTGTTCCA AATTAACGT 2880
TTAACCTTT TAATAGGTTT ATACAATAAT TGATATATGT TTCTGTATA TGTCTAATT 2940
GTTATCATCC ATTTAGATAT AGACGAAAAA AAATCTAAGA ACTAAAAAAT ATGCTAATTT 3000
35 GAAATGAAGG GAGTATATAT TGGGATAATG TCGATGAGAT CCCTCGTAAT ATCACCAGCA 3060
TCACACGTGT CCAGTTAATG TATCAGTGAT ACGTGTATTG ACATTTGTTG CGCGTAGGCG 3120
TACCAACAA TTTTGATCGA CTATCAGAAA GTCAACGGAA GCGAGTCGAC CTCGAGGGGG 3180
GGCCGGTAC CCAGCTTTTG TTCCCTTTAG TGAGGGTTAA TTGCGCGCTT GGCCTAATCA 3240
TGGTCTATAG TGTTCCTGT GTGAAATTGT TATCCGCTCA CAATTCCACA CAACATACGA 3300
40 GCCGGAAGCA TAAAGTGTA AGCCTGGGGT GCCTAATGAG TGAGCTAACT CACATTAATT 3360
GCGTTGCGCT CACTGCCCGC TTTCCAGTCG GGAACCTGT CGTGCCAGCT GCATTAATGA 3420
ATCGGCCAAC GCGCGGGGAG AGGCGGTTTG CGTATTGGGC GCTCTCCGC TTCCTCGCTC 3480
ACTGACTCGC TGCGCTCGGT CGTTCCGGTG CGGCGAGCGG TATCAGCTCA CTCAAAGGCG 3540
GTAATACGGT TATCCACAGA ATCAGGGGAT AACGACGAA AGAACATGTG AGCAAAAGGC 3600
45 CAGCAAAAGG CCAGGAACCG TAAAAAGGCC CGGTTGCTGG CGTTTTTCCA TAGGCTCCG 3660
CCCCCTGACG AGCATCACA AAATCGACGC TCAAGTCAGA GGTGGCGAAA CCGACAGGA 3720
CTATAAAGAT ACCAGGCGTT TCCCCCTGGA AGCTCCCTCG TCGCTCTCC TGTTCCGACC 3780
CTGCCGCTTA CCGGATACCT GTCCGCTTTT CTCCCTTCGG GAAGCGTGGC GCTTCTCAT 3840
AGCTCACGCT GTAGGTATCT CAGTTCGGTG TAGGTGTTT GCTCCAAGCT GGGCTGTGTG 3900
50 CACGAACCCC CGTTTCAGCC CGACCGTTGC GCCTATCCG GTAACATATCG TCTTGAGTCC 3960
AACCCGGTAA GACACGACTT ATCGCCACTG CGACGACCA CTGGTAACAG GATTAGCAGA 4020
GCGAGGTATG TAGGCGGTGC TACAGAGTTC TTGAAGTGGT GGCCTAACTA CGGCTACACT 4080
AGAAGGACAG TATTTGGTAT CTGCGCTCTG CTGAAGCCAG TTACCTTCGG AAAAAGAGTT 4140
GGTAGCTCTT GATCCGGCAA ACAAACCACC GCTGGTAGCG GTGGTTTTTT TGTTTGCAAG 4200
55 CAGCAGATTA CGCGCAGAAA AAAAGGATCT CAAGAAGATC CTTTGATCTT TTCTACGGGG 4260
TCTGACGCTC AGTGAACGA AAACCTACGT TAAGGGATTG TGGTCATGAG ATTATCAAAA 4320
AGGATCTTCA CCTAGATCCT TTAAATTA AAATGAAGTT TAAATCAAT CTAAGTATA 4380
TATGAGTAAA CTTGGTCTGA CAGTTACCAA TGCTTAATCA GTGAGGCACC TATCTCAGCG 4440
ATCTGTCTAT TTCGTTTATC CATAGTTGCC TGACTCCCCG TCGTGTAGAT AACTACGATA 4500
60 CGGGAGGGCT TACCATCTGG CCGAGTGCT GCAATGATAC CGCGAGACCC ACGCTCACCG 4560
GCTCCAGATT TATCAGCAAT AAACCAGCCA GCCGGAAGGG CCGAGCGCAG AAGTGGTCTT 4620
GCAACTTTAT CCGCTCCAT CCAGTCTATT AATTGTTGCC GGGAAGCTAG AGTAAGTAGT 4680
TCGCCAGTTA ATAGTTTGGC CAACGTTGTT GCCATTGCTA CAGGCATCGT GGTGTACAGC 4740
TCGTCGTTTG GTATGGCTTC ATTCAGCTCC GGTCCCAAC GATCAAGGCG AGTTACATGA 4800
65 TCCCCATGT TGTGCAAAAA AGCGGTTAGC TCCTTCGGTC CTCCGATCGT TGTGAGAAGT 4860
AAGTTGGCCG CAGTGTTATC ACTCATGGTT ATGGCAGCAC TGCATAATTC TCTTACTGTC 4920

ATGCCATCCG TAAGATGCTT TTCTGTGACT GGTGAGTACT CAACCAAGTC ATTCTGAGAA 4980
TAGTGTATGC GGGCACCAGG TTGCTCTTGC CCGGCGTCAA TACGGGATAA TACCGCGCCA 5040
CATAGCAGAA CTTTAAAAGT GCTCATCATT GGAAAACGTT CTTGCGGGCG AAAACTCTCA 5100
AGGATCTTAC CGCTGTTGAG ATCCAGTTCG ATGTAACCCA CTCGTGCACC CAACTGATCT 5160
5 TCAGCATCTT TTACTTTTAC CAGCGTTTCT GGGTGAGCAA AAACAGGAAG GCAAAATGCC 5220
GCAAAAAAGG GAATAAGGGC GACACGGAAA TGTTGAATAC TCATACTCTT CCTTTTCAA 5280
TATTATTGAA GCATTTATCA GGGTTATTGT CTCATGAGCG GATACATATT TGAATGTATT 5340
TAGAAAAATA AACAAATAGG GGTTCGCGC ACATTTCCCC GAAAAGTGCC AC 5392

10 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5173 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CTAAATTGTA AGCGTTAATA TTTTGTTAAA ATTCGCGTTA AATTTTGT AAATCAGCTC 60
ATTTTAAAC CAATAGGCCG AAATCGGCAA AATCCCTTAT AAATCAAAAG AATAGACCGA 120
GATAGGGTTG AGTGTGTTC CAGTTTGGA CAAGAGTCCA CTATTAAAGA ACGTGGACTC 180
25 CAACGTCAA GGGCGAAAA CCGTCTATCA GGGCGATGGC CCACTACGTG AACCATCACC 240
CTAATCAAGT TTTTGGGGT CGAGGTGCCG TAAAGCACTA AATCGGAACC CTAAAGGGAG 300
CCCCGATTT AGAGCTTGAC GGGGAAAGCC GCGAACGTG GCGAGAAAGG AAGGGAAGAA 360
AGCGAAAGGA GCGGGCGCTA GGGCGCTGGC AAGTGTAGCG GTCACGCTGC GCGTAACCA 420
CACACCCGCC GCGCTTAATG CGCCGCTACA GGGCGGTCC CATTGCCAT TCAGGCTGCG 480
30 CAACTGTTGG GAAGGGCGAT CGGTGCGGGC CTCTTCGCTA TTACGCCAGC TGGCGAAAGG 540
GGGATGTGCT GCAAGGCGAT TAAGTTGGGT AACGCCAGGG TTTTCCAGT CACGACGTTG 600
TAAACGACG GCCAGTGAGC GCGCGTAATA CGACTCACTA TAGGGCGAAT TGGAGCTCCA 660
CCGCGGTGGC GGCCGCTCTA GATTATATAA TTTATAAGCT AAACAACCCG GCCCTAAAGC 720
ACTATCGTAT CACCTATCTA AATAAGTCAC GGGAGTTTCG AACGTCCACT TCGTCGCACG 780
35 GAATTGCATG TTTCTTGTG GAAGCATATT CACGCAATCT CCACACATAA AGGTTTATGT 840
ATAAACTTAC ATTTAGCTCA GTTTAATTAC AGTCTTATTT GGATGCATAT GTATGGTTCT 900
CAATCCATAT AAGTTAGAGT AAAAAATAAG TTAAATTTT ATCTTAATTC ACTCCAACAT 960
ATATGGATCT ACAATACTCA TGTGCATCCA AACAACTAC TTATATTGAG GTGAATTTGG 1020
TAGAAATTAA ACTAATTAC ACTAAGCC AATCTTACT ATATTAAAGC ACCAGTTTCA 1080
40 ACGATCGTCC CGCGTCAATA TTATTAAAA ACTCCTACAT TTCTTTATAA TCAACCCGCA 1140
CTCTTATAAT CTCTTCTCTA CTACTATAA AAGAGAGTTT ATGTACAAAA TAAGGTGAAA 1200
TTATCTATAA GTGTTCTGGA TATTGGTTGT TGGCTCCCAT ATTCACACAA CCTAATCAAT 1260
AGAAAACATA TGTTTTATTA AAACAAATT TATCATATAT CATATATATA TATATATCAT 1320
ATATATATAT AAACCGTAGC AATGCACGGG CATATACTA GTGCAACTTA ATACATGTGT 1380
45 GTATTAAGAT GAATAAGAGG GTATCCAAAT AAAAACTTG TTGCTTACGT ATGGATCGAA 1440
AGGGGTTGGA AACGATTAAA CGATTAAATC TCTTCCTAGT CAAAATTGAA TAGAAGGAGA 1500
TTTAATATAT CCCAATCCCC TTCGATCATC CAGGTGCAAC CGTATAAGTC CTAAGTGGT 1560
GAGGAACACG AAAGAACCAT GCATTGGCAT GTAAAGCTCC AAGAATTTGT TGTATCTTCA 1620
ACAACCTACA GAACATCAAC CAAAATTGCA CGTCAAGGGT ATTGGGTAAG AAACAATCAA 1680
50 ACAATCCTC TCTGTGTGCA AAGAAACACG GTGAGTCATG CCGAGATCAT ACTCATCTGA 1740
TATACATGCT TACAGCTCAC AAGACATTAC AAACAACCTA TATTGCATTA CAAAGATCGT 1800
TTCATGAAAA ATAAATAGG CCGGACAGGA CAAAATCCT TGACGTGTAA AGTAAATTTA 1860
CAACAAAAAA AAAGCCATAT GTCAAGCTAA ATCTAATTCG TTTTACGTAG ATCAACAACC 1920
TGTATAAGG AACAAAACTG AGCCAGCAG AAGTACAGAA TGATTCCAGA TGAACCATCG 1980
55 ACGTGTACG TAAAGAGAGT AAGAGTCAT ATACATTTGG CAAGAAACCA TGAAGCTGCC 2040
TACAGCCGTA TCGGTGGCAT AAGAACACAA GAAATTGTGT TAATTAATCA AAGCTATAAA 2100
TAACGCTCGC ATGCTGTGC ACTTCTCCAT CACCACCACT GGGTCTTCAG ACCATTAGCT 2160
TTATCTACTC CAGAGCGCAG AAGAACCCGA TCGACACCAT GAAGTCGGTG GAGAAGAAAC 2220
CGAAGGGTGT GAAGACAGGT GCGGGTGACA AGCATAAGCT GAAGACAGAG TGGCCGGAGT 2280
60 TGGTGGGGAA ATCGGTGGAG AAAGCCAAGA AGGTGATCCT GAAGGACAAG CCAGAGGCGC 2340
AAATCATAGT TCTACCGGTT GGTACAAAGG TGGGTAAGCA TTATAAGATC GACAAGGTCA 2400
AGCTTTTGT GGATAAAAAA GACAACATCG CGCAGGTCCC CAGGGTCGGC TAGCCTCGAG 2460
ATCCCCGGCG GTGTCCCCCA CTGAAGAAAC TATGTGCTGT AGTATAGCCG CTGGCTAGCT 2520
AGCTAGTTGA GTCATTTAGC GGCGATGATT GAGTAATAAT GTGTCACGCA TCACCATGCA 2580
65 TGGGTGGCAG TCTCAGTGTG AGCAATGACC TGAATGAACA ATTGAAATGA AAAGAAAAAA 2640
GTATTGTTCC AAATTAAACG TTTTAACCTT TTAATAGGTT TATACAATAA TTGATATATG 2700

TTTTCTGTAT ATGTCTAATT TGTATCATC CATTTAGATA TAGACGAAAA AAAATCTAAG 2760
AACTAAAACA AATGCTAATT TGAAATGAAG GGAGTATATA TTGGGATAAT GTCGATGAGA 2820
TCCCTCGTAA TATCACCGAC ATCACACGTG TCCAGTTAAT GTATCAGTGA TACGTGTATT 2880
CACATITGTT GCGCGTAGGC GTACCCAACA ATTTTGATCG ACTATCAGAA AGTCAACGGA 2940
5 AGCGAGTCGA CCTCGAGGGG GGGCCCGGTA CCCAGCTTTT GTTCCCTTTA GTGAGGGTTA 3000
ATTGCGCGCT TGGCGTAATC ATGGTCATAG CTGTTTCCTG TGTGAAATTG TTATCCGCTC 3060
ACAATTCCAC ACAACATACG AGCCGGAAGC ATAAAGTGTA AAGCCTGGGG TGCCTAATGA 3120
GTGAGCTAAC TCACATTAAT TGC GTTGC GC TCACTGCCCG CTTTCCAGTC GGGAAACCTG 3180
TCGTGCCAGC TGCATTAATG AATCGGCCAA CGCGCGGGGA GAGGCGGTTT GCGTATTGGG 3240
10 CGCTCTTCCG CTTCCTCGCT CACTGACTCG CTGCGCTCGG TCGTTCGGCT GCGGCGAGCG 3300
GTATCAGCTC ACTCAAAGGC GGTAAACGG TTATCCACAG AATCAGGGGA TAACGCAGGA 3360
AAGAACATGT GAGCAAAAGG CCAGCAAAAG GCCAGGAACC GTAAAAAGGC CGCGTTGCTG 3420
GCGTTTTTCC ATAGGCTCCG CCCCCGTGAC GAGCATCACA AAAATCGACG CTCAAGTCAG 3480
AGGTGGCGAA ACCCGACAGG ACTATAAAGA TACCAGGCGT TTCCCCCTGG AAGCTCCCTC 3540
15 GTGCGCTCTC CTGTTCCGAC CCTGCCGCTT ACCGGATACC TGTCCGCCTT TCTCCCTTCG 3600
GGAAGCGTGG CGCTTTCTCA TAGCTCACGC TGTAGGTATC TCAGTTCGGT GTAGGTCGTT 3660
CGTCCAAGC TGGGCTGTGT GCACGAACCC CCCGTTGAGC CCGACCGCTG CGCCTTATCC 3720
GGTAACTATC GTCTTGAGTC CAACCCGGTA AGACACGACT TATCGCCACT GGCAGCAGCC 3780
ACTGGTAACA GGATTAGCAG AGCGAGGTAT GTAGGCGGTG CTACAGAGTT CTTGAAGTGG 3840
20 TGGCCTAACT ACGGCTACAC TAGAAGGACA GTATTTGGTA TCTGCGCTCT GCTGAAGCCA 3900
GTTACCTTCG GAAAAAGAGT TGGTAGCTCT TGATCCGGCA AACAAACCAC CGCTGGTAGC 3960
GGTGGTTTTT TTGTTTGCAA GCAGCAGATT ACGCGCAGAA AAAAAAGGATC TCAAGAAGAT 4020
CCTTTGATCT TTTCTACGGG GTCTGACGCT CAGTGGAACG AAAACTCACG TTAAGGGATT 4080
TTGGTCATGA GATTCTGAAA AAGGATCTTC ACCTAGATCC TTTTAAATTA AAAATGAAGT 4140
25 TTTAAATCAA TCTAAAGTAT ATATGAGTAA ACTTGGTCTG ACAGTTACCA ATGCTTAATC 4200
AGTGAGGCAC CTATCTCAGC GATCTGTCTA TTTCGTTTCA CCATAGTTGC CTGACTCCCC 4260
GTCGTGTAGA TAACTACGAT ACGGGAGGGG TTACCATCTG GCCCCAGTGC TGCAATGATA 4320
CCGCGAGACC CAGCTCACC GGCTCCAGAT TTATCAGCAA TAAACCAGCC AGCCGGAAGG 4380
GCCGAGCGCA GAGTGGTCC TGCAACTTTA TCCGCCTCCA TCCAGTCTAT TAATTGTTGC 4440
30 CGGGAAGCTA GAGTAAGTAG TTCGCCAGTT AATAGTTTGC GCAACGTTGT TGCCATTGCT 4500
ACAGGCATCG TGGTGTACG CTCGTCGTTT GGTATGGCTT CATTGAGTCT CGGTTCCCAA 4560
CGATCAAGGC GAGTTACATG ATCCCCCATG TTGTGCAAAA AAGCGGTTAG CTCCTTCGGT 4620
CCTCCGATCG TTGTCAGAAG TAAGTTGGCC GCAGTGTTAT CACTCATGGT TATGGCAGCA 4680
CTGCATAATT CTCTTACTGT CATGCCATCC GTAAGATGCT TTTCTGTGAC TGGTGAGTAC 4740
35 TCAACCAAGT CATTCTGAGA ATAGTGATG CGGCGACCGA GTTGCTCTTG CCCGGCGTCA 4800
ATACGGGATA ATACCGCGCC ACATAGCAGA ACTTTAAAAG TGCTCATCAT TGGAAAACGT 4860
TCTTCGGGGC GAAAACTCTC AAGGATCTTA CCGCTGTTGA GATCCAGTTC GATGTAACCC 4920
ACTCGTGAC CCAACTGATC TTCAGCATCT TTTACTTTCA CCAGCGTTTC TGGGTGAGCA 4980
AAAACAGGAA GGCAAAATGC CGCAAAAAAG GGAATAAGGG CGACACGGAA ATGTTGAATA 5040
40 CTCATACCTT TCTTTTCA ATATTATTGA AGCATTTATC AGGGTTATTG TCTCATGAGC 5100
GGATACATAT TTGAATGTAT TTAGAAAAAT AAACAAATAG GGGTTCGCG CACATTTCCC 5160
CGAAAAGTGC CAC 5173

(2) INFORMATION FOR SEQ ID NO:8:

45

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

55

AGTATAAGTA AACACACCAT CACACCCTTG AGGCCCTTGC TGGTGGCCAT GGTG 54

(2) INFORMATION FOR SEQ ID NO:9:

60

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
65 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

5 CCTCACATCC CTTAGTGCCT AAGTTCGACG TCGGGCCCTC TAGTCGACGG ATCCA 55

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

AGCGGAAAAT GCCCGAAAGG CTTCCCCAAA TTGGC 35

20 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TGCGCAGGCG TCTGCAAGTG TAAGCTGACT AGTAGCGGAA AATGC 45

(2) INFORMATION FOR SEQ ID NO:12:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

45 TACAACCTTT GCAAAGTCAA AGGCGCCAAG AAGCTTTGCG CAGGCGTCTG 50

(2) INFORMATION FOR SEQ ID NO:13:

50 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

60 GCAAGAGTTG CTGCAAGAGT ACCCTGGGAA GGAAGTGCTA CAACCTTTGC 50

INTERNATIONAL SEARCH REPORT

International Application No

PCT/99/02061

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/82 C12N15/29 A01H5/00 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 28247 A (BIOCEM ;LUDEVID DOLORES (ES); TORRENT MARGARITA (ES); ALVAREZ INAK) 7 August 1997 (1997-08-07)	1-3,5-7, 9-21
Y	the whole document	4,8,14
Y	WO 93 03160 A (DU PONT) 18 February 1993 (1993-02-18) page 92, line 8 - line 10	4
Y	WO 97 35023 A (PIONEER HI BRED INT) 25 September 1997 (1997-09-25) page 19, line 6 - line 9	8,14
Y	WO 97 41239 A (PIONEER HI BRED INT ;BEACH LARRY R (US)) 6 November 1997 (1997-11-06) the whole document	8

	---/---	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 July 1999

Date of mailing of the international search report

14/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/02061

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94 16078 A (PIONEER HI BRED INT) 21 July 1994 (1994-07-21) cited in the application the whole document ---	8
Y	WO 96 38562 A (PIONEER HI BRED INT) 5 December 1996 (1996-12-05) cited in the application the whole document ---	8
Y	WO 96 38563 A (PIONEER HI BRED INT) 5 December 1996 (1996-12-05) cited in the application the whole document ---	8
X	WO 98 02563 A (PIONEER HI BRED INT ; BEACH LARRY (US); COLEMAN CRAIG E (US); LARKI) 22 January 1998 (1998-01-22) page 9, line 6 - line 28 page 11, line 13 - line 25; claims 1-7 ---	1-13, 15-21
X	WO 95 06128 A (DEKALB GENETICS CORP) 2 March 1995 (1995-03-02) page 211 - page 222 ---	1,5-7, 9-18,20, 21
X	WO 97 26366 A (DEKALB GENETICS CORP ; ANDERSON PAUL C (US); CHOMET PAUL S (US); GR) 24 July 1997 (1997-07-24) page 67 ---	1,2,5,6, 9-12
X	WO 93 08682 A (STATE UNIVERSITY OF NEW JERSEY) 13 May 1993 (1993-05-13) the whole document ---	1,5-7, 9-12
X	WO 91 10725 A (BIOTEKNIKA INTERNATIONAL) 25 July 1991 (1991-07-25) page 14, line 35 - page 16, line 16 page 44, line 1 - page 56, line 26: claims 1-7 ---	13-17
X	WO 96 38574 A (PIONEER HI BRED INT) 5 December 1996 (1996-12-05) page 10, line 34 - page 16, line 14 ---	13-17
A	WO 98 26064 A (DEKALB GENETICS CORP) 18 June 1998 (1998-06-18) examples 8,9 ---	1
P,X	WO 98 26064 A (DEKALB GENETICS CORP) 18 June 1998 (1998-06-18) examples 8,9 ---	1-7,9-21

	-/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT 99/02061

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TORRENT M ET AL: "LYSINE-RICH MODIFIED GAMMA-ZEINS ACCUMULATE IN PROTEIN BODIES OF TRANSIENTLY TRANSFORMED MAIZE ENDOSPERMS" PLANT MOLECULAR BIOLOGY, vol. 34, no. 1, May 1997 (1997-05), pages 139-149, XP002034740 ISSN: 0167-4412 the whole document ---	1-21
A	WO 95 15392 A (DU PONT ;FALCO SAVERIO CARL (US); KEELER SHARON JO (US); RICE JANE) 8 June 1995 (1995-06-08) examples 12,13 ---	1-21
A	WO 96 01905 A (DU PONT ;FALCO SAVERIO CARL (US)) 25 January 1996 (1996-01-25) the whole document ---	2
A	WO 89 04371 A (UNIV LOUISIANA STATE) 18 May 1989 (1989-05-18) example 15 ---	1,5,19
E	WO 99 04024 A (DOW AGROSCIENCES LLC) 28 January 1999 (1999-01-28) examples 1-11 -----	1-3,5-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/02061

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9728247 A	07-08-1997	FR 2744134 A AU 1550297 A CA 2242903 A EP 0877754 A	01-08-1997 22-08-1997 07-08-1997 18-11-1998
WO 9303160 A	18-02-1993	AU 661334 B AU 2441292 A CA 2114788 A EP 0598806 A JP 7502163 T US 5559223 A ZA 9205984 A	20-07-1995 02-03-1993 18-02-1993 01-06-1994 09-03-1995 24-09-1996 10-02-1994
WO 9735023 A	25-09-1997	US 5850016 A AU 2535197 A CA 2220959 A CN 1193355 A EP 0828846 A PL 323627 A	15-12-1998 10-10-1997 25-09-1997 16-09-1998 18-03-1998 14-04-1998
WO 9741239 A	06-11-1997	AU 2729597 A	19-11-1997
WO 9416078 A	21-07-1994	AU 6162294 A CA 2161881 A EP 0745126 A	15-08-1994 21-07-1994 04-12-1996
WO 9638562 A	05-12-1996	AU 705933 B AU 5961096 A CA 2222615 A CN 1192238 A EP 0828835 A PL 323641 A US 5885801 A	03-06-1999 18-12-1996 05-12-1996 02-09-1998 18-03-1998 14-04-1998 23-03-1999
WO 9638563 A	05-12-1996	AU 5961196 A CA 2222600 A CN 1192239 A EP 0832235 A PL 323635 A US 5885802 A	18-12-1996 05-12-1996 02-09-1998 01-04-1998 14-04-1998 23-03-1999
WO 9802563 A	22-01-1998	CA 2181418 A AU 3651697 A	18-01-1998 09-02-1998
WO 9506128 A	02-03-1995	AU 5640498 A AU 684105 B AU 7716994 A BR 9407355 A CA 2170260 A EP 0721509 A HU 74392 A US 5780709 A ZA 9406488 A ZA 9604217 A	04-06-1998 04-12-1997 21-03-1995 19-08-1997 02-03-1995 17-07-1996 30-12-1996 14-07-1998 30-11-1995 26-08-1996
WO 9726366 A	24-07-1997	AU 1706697 A CA 2243268 A EP 0879292 A	11-08-1997 24-07-1997 25-11-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No
PCT/99/02061

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9308682 A	13-05-1993	NONE	
WO 9110725 A	25-07-1991	US 5484956 A CA 2074355 A CN 1054170 A US 5538880 A US 5508468 A US 5554798 A US 5780708 A US 5538877 A	16-01-1996 23-07-1991 04-09-1991 23-07-1996 16-04-1996 10-09-1996 14-07-1998 23-07-1996
WO 9638574 A	05-12-1996	AU 5956196 A CA 2222673 A CN 1195377 A EP 0828845 A	18-12-1996 05-12-1996 07-10-1998 18-03-1998
WO 9826064 A	18-06-1998	AU 5795898 A	03-07-1998
WO 9515392 A	08-06-1995	AU 704510 B AU 1255995 A BR 9408228 A CA 2177351 A EP 0734445 A HU 75078 A JP 9511124 T PL 314696 A ZA 9409385 A	22-04-1999 19-06-1995 26-08-1997 08-06-1995 02-10-1996 28-04-1997 11-11-1997 16-09-1996 27-05-1996
WO 9601905 A	25-01-1996	AU 2963695 A BR 9510174 A CA 2192550 A EP 0769061 A HU 77112 A	09-02-1996 04-11-1997 25-01-1996 23-04-1997 02-03-1998
WO 8904371 A	18-05-1989	AU 2802989 A CA 1321157 A EP 0675960 A US 5597945 A US 5811654 A	01-06-1989 10-08-1993 11-10-1995 28-01-1997 22-09-1998
WO 9904024 A	28-01-1999	AU 8148398 A	10-02-1999

